

DUALIZABLE ALGEBRAS WITH PARALLELOGRAM TERMS

KEITH A. KEARNES AND ÁGNES SZENDREI

ABSTRACT. We prove that if $\mathbf{A}$ is a finite algebra with a parallelogram term that satisfies the split centralizer condition, then $\mathbf{A}$ is dualizable. This yields yet another proof of the dualizability of any finite algebra with a near unanimity term, but more importantly proves that every finite module, group or ring in a residually small variety is dualizable.

1. INTRODUCTION

Natural Duality Theory investigates categorical dualities that are mediated by finite algebras. One of the main goals of the theory is to identify those finite algebras that can serve as character algebras for a duality. Such algebras are called *dualizable*. Although general criteria for dualizability are known, the problem of identifying the finite algebras satisfying the criteria is still difficult. The broadest natural class of algebras where the problem has been solved is the class of finite algebras $\mathbf{A}$ such that the prevariety $\mathbf{SP}(\mathbf{A})$ has tame congruence-theoretic typeset contained in $\{\mathbf{3}, \mathbf{4}\}$. According to the Big NU Obstacle Theorem from [5] a finite algebra satisfying this assumption is dualizable if and only if it has a near unanimity term operation.

This paper probes the broader class of finite algebras $\mathbf{A}$ such that the prevariety $\mathbf{SP}(\mathbf{A})$ has tame congruence-theoretic typeset contained in $\{\mathbf{2}, \mathbf{3}, \mathbf{4}\}$. There is a natural analogue of a near unanimity term operation that is likely to be involved in this context, called a “parallelogram term operation”. Recently M. Moore [16] has announced an extension of one direction of the Big NU Obstacle Theorem, namely, under the assumption that $\mathbf{A}$ is finite and $\mathbf{SP}(\mathbf{A})$ has tame congruence-theoretic typeset contained in $\{\mathbf{2}, \mathbf{3}, \mathbf{4}\}$, it is the case that if $\mathbf{A}$ is dualizable then it must have a parallelogram term operation. Known examples show that an unmodified converse to this statement cannot hold. Thus, we expect that there is some condition ε such that, if $\mathbf{A}$ is finite and $\mathbf{SP}(\mathbf{A})$ has tame congruence-theoretic typeset contained in $\{\mathbf{2}, \mathbf{3}, \mathbf{4}\}$, then $\mathbf{A}$ is dualizable if and only if $\mathbf{A}$ has a parallelogram term operation and condition ε holds.

2010 *Mathematics Subject Classification.* Primary: 08C20; Secondary: 08A05, 08A40.

Key words and phrases. natural duality, dualizable algebra, parallelogram term, cube term.

This material is based upon work supported by the Hungarian National Foundation for Scientific Research (OTKA) grant no. K83219 and K104251.

In this paper we produce a candidate for condition ε under the additional assumption that $\mathbf{A}$ lies in a residually small variety. We call the condition the “split centralizer condition”. Specifically, we operate under the global assumptions that $\mathbf{A}$ is a finite algebra from a residually small variety and $\mathbf{SP}(\mathbf{A})$ has tame congruence-theoretic typeset contained in $\{\mathbf{2}, \mathbf{3}, \mathbf{4}\}$. We conjecture that, under the global assumptions, $\mathbf{A}$ will be dualizable if and only if $\mathbf{A}$ has a parallelogram term operation and satisfies the split centralizer condition. What we prove in this paper is the “if” part of this conjecture.

Let us reformulate the statement that we prove. It is a fact that a finite algebra with a parallelogram term automatically has the property that $\mathbf{SP}(\mathbf{A})$ has tame congruence-theoretic typeset contained in $\{\mathbf{2}, \mathbf{3}, \mathbf{4}\}$. Thus, our result is that if $\mathbf{A}$ is finite, lies in a residually small variety, has a parallelogram term, and satisfies the split centralizer condition, then $\mathbf{A}$ is dualizable. Interestingly, our result is strong enough to prove the dualizability of the dualizable algebras in residually small varieties with a parallelogram term in all the known cases, and in many new cases, and yet neatly avoids an obstacle to dualizability identified by Idziak in 1994.

Let us state the split centralizer condition. Let $\mathbf{B}$ be a finite algebra and let $\mathcal{Q}$ be a quasivariety containing $\mathbf{B}$. A $\mathcal{Q}$ -congruence on $\mathbf{B}$ is a congruence κ of $\mathbf{B}$ such that $\mathbf{B}/\kappa \in \mathcal{Q}$. Let δ be a meet irreducible congruence on $\mathbf{B}$ with upper cover θ , and let $\nu = (\delta : \theta)$ be the centralizer of θ modulo δ . We will say that the triple of congruences (δ, θ, ν) which arises in this way is *split* (relative to $\mathcal{Q}$) by a triple (α, β, κ) of congruences of $\mathbf{B}$ if

- (i) κ is a $\mathcal{Q}$ -congruence on $\mathbf{B}$,
- (ii) $\beta \leq \delta$,
- (iii) $\alpha \wedge \beta = \kappa$,
- (iv) $\alpha \vee \beta = \nu$, and
- (v) α/κ is abelian (or equivalently, $[\alpha, \alpha] \leq \kappa$).

Now, if θ/δ is nonabelian, then $\nu = (\delta : \theta) = \delta$, so (δ, θ, ν) will be split by $(\alpha, \beta, \kappa) := (0, \delta, 0)$. Therefore splitting is only in question when θ/δ is abelian. Henceforth we shall focus only on this case and call a triple (δ, θ, ν) of congruences on $\mathbf{B}$ *relevant* if

- (a) δ is completely meet irreducible,
- (b) θ is the upper cover of δ ,
- (c) $\nu = (\delta : \theta)$, and
- (d) θ/δ is abelian.

The *split centralizer condition* for a finite algebra $\mathbf{A}$ is the condition that, for $\mathcal{Q} := \mathbf{SP}(\mathbf{A})$ and for any subalgebra $\mathbf{B} \leq \mathbf{A}$, each relevant triple (δ, θ, ν) of $\mathbf{B}$ is split (relative to $\mathcal{Q}$) by some triple (α, β, κ) . The relationships between these congruences of $\mathbf{B}$ is depicted in Figure 1.




FIGURE 1.

It is not hard to show that if a finite algebra with a parallelogram term satisfies the split centralizer condition, then it generates a residually small variety (combine Corollary 2.2 and Theorem 2.5). Therefore our main result can be restated as follows.

Theorem 1.1. *If $\mathbf{A}$ is a finite algebra with a parallelogram term and $\mathbf{A}$ satisfies the split centralizer condition, then $\mathbf{A}$ is dualizable.*

Section 2 summarizes preliminaries. Sections 3 through 5 are devoted to proving Theorem 1.1. In Section 6, we apply Theorem 1.1 to prove that if $\mathcal{V}$ is a variety with a parallelogram term in which every finite subdirectly irreducible algebra is either abelian or neutral or almost neutral, then every finite algebra in $\mathcal{V}$ is dualizable (Theorem 6.4). This generalizes the following known results:

- (1) every finite algebra with a near unanimity term is dualizable [7],
- (2) every paraprimal algebra is dualizable [6], and
- (3) every finite commutative ring R with unity whose Jacobson radical squares to zero is dualizable [3].

Theorem 6.4 also implies the new results that

- (4) every finite module is dualizable, and
- (5) every finite ring (commutative or not, unital or not) in a residually small variety is dualizable.

We also show how to apply Theorem 1.1 to deduce that every finite group with abelian Sylow subgroups is dualizable [17]. This last theorem is not a consequence of Theorem 6.4.

2. PRELIMINARIES

Algebras will be denoted by boldface letters, their universes by the same letters in italics. For arbitrary algebras $\mathbf{A}$ and $\mathbf{B}$, $\text{Aut}(\mathbf{A})$ denotes the automorphism group of

$\mathbf{A}$, $\text{Con}(\mathbf{A})$ the congruence lattice of $\mathbf{A}$, and $\text{Hom}(\mathbf{A}, \mathbf{B})$ the set of all homomorphisms $\mathbf{A} \rightarrow \mathbf{B}$. The top and bottom elements of $\text{Con}(\mathbf{A})$ are denoted 1 and 0, respectively, and the identity map on any set A is denoted id_A . The variety generated by an algebra $\mathbf{A}$ is denoted by $\mathcal{V}(\mathbf{A})$. We will write $\mathbf{B} \leq \mathbf{A}$ to indicate that $\mathbf{B}$ is a subalgebra of $\mathbf{A}$. By a *section* of $\mathbf{A}$ we mean a quotient of a subalgebra of $\mathbf{A}$.

Let ϑ be an equivalence relation on a set A . The ϑ -class of an element $a \in A$ is denoted by a/ϑ , and the number of equivalence classes of ϑ may be referred to as *the index of ϑ* . We will often write $a \equiv_{\vartheta} b$ instead of $(a, b) \in \vartheta$. For $B \subseteq A$ the restriction of ϑ to B is denoted by ϑ_B . If $\mathbf{B} \leq \mathbf{A}$ and ϑ is a congruence of $\mathbf{A}$, we may write $\vartheta_{\mathbf{B}}$ for ϑ_B , which is a congruence of $\mathbf{B}$.

Let ϑ be a congruence of an algebra $\mathbf{A}$, and let $\mathbf{B}$ be a subalgebra of $\mathbf{A}$. We say that $\mathbf{B}$ is *saturated with respect to ϑ* , or $\mathbf{B}$ is a *ϑ -saturated subalgebra* of $\mathbf{A}$, if $b \in \mathbf{B}$ and $b \equiv_{\vartheta} a$ imply $a \in \mathbf{B}$ for all $a \in \mathbf{A}$. In other words, $\mathbf{B}$ is ϑ -saturated if and only if its universe is a union of ϑ -classes of $\mathbf{A}$. For arbitrary subalgebra $\mathbf{B}$ of $\mathbf{A}$ there exists a smallest ϑ -saturated subalgebra of $\mathbf{A}$ that contains $\mathbf{B}$, which we denote by $\mathbf{B}[\vartheta]$; the universe of $\mathbf{B}[\vartheta]$ is $B[\vartheta] := \bigcup_{b \in B} b/\vartheta$. By the second isomorphism theorem the map $\mathbf{B}/\vartheta_{\mathbf{B}} \rightarrow \mathbf{B}[\vartheta]/\vartheta_{\mathbf{B}[\vartheta]}$, $b/\vartheta_{\mathbf{B}} \mapsto b/\vartheta_{\mathbf{B}[\vartheta]} (= b/\vartheta)$ is an isomorphism, so the indices of $\vartheta_{\mathbf{B}}$ and $\vartheta_{\mathbf{B}[\vartheta]}$ are equal.

For every natural number m we will use the notation $[m]$ for the set $\{1, 2, \dots, m\}$.

2.1. The Modular Commutator and Residual Smallness. For the definition and basic properties of the commutator operation $[\ , \]$ on congruence lattices of algebras in congruence modular varieties the reader is referred to [10]. To avoid confusion, intervals in congruence lattices will be denoted by $\llbracket \ , \ \rrbracket$. Recall that a congruence $\alpha \in \text{Con}(\mathbf{A})$ of an algebra $\mathbf{A}$ is called *abelian* if $[\alpha, \alpha] = 0$, and an interval $\llbracket \beta, \alpha \rrbracket$ in $\text{Con}(\mathbf{A})$ is called abelian if $[\alpha, \alpha] \leq \beta$, or equivalently, if the congruence $\alpha/\beta \in \text{Con}(\mathbf{A}/\beta)$ is abelian. For two congruences $\beta \leq \alpha$ in $\text{Con}(\mathbf{A})$ the *centralizer of α modulo β* , denoted $(\beta : \alpha)$, is the largest congruence $\gamma \in \text{Con}(\mathbf{A})$ such that $[\alpha, \gamma] \leq \beta$. The *center* of $\mathbf{A}$ is the congruence $\zeta := (0 : 1)$.

For a cardinal c , a variety $\mathcal{V}$ is called *residually less than c* if every subdirectly irreducible algebra in $\mathcal{V}$ has cardinality $< c$; $\mathcal{V}$ is called *residually small* if it is residually less than some cardinal. It is proved in [10, Theorem 10.14] that for a congruence modular variety $\mathcal{V}$ to be residually small, it is necessary that the congruence lattice of every algebra $\mathbf{A} \in \mathcal{V}$ satisfy the commutator identity

$$(C1) \quad [x \wedge y, y] = x \wedge [y, y],$$

which can also be expressed by the implication $x \leq [y, y] \rightarrow x = [x, y]$ (see [10, Theorem 8.1]). For finitely generated varieties the converse is also true, as the following theorem states.

Theorem 2.1 (From [10, Theorem 10.15]). *Let $\mathbf{A}$ be a finite algebra that generates a congruence modular variety $\mathcal{V}(\mathbf{A})$. Then the following conditions are equivalent:*

- (a) $\mathcal{V}(\mathbf{A})$ is residually small,
- (b) $\mathcal{V}(\mathbf{A})$ is residually $< q$ for some natural number q ,
- (c) (C1) holds in the congruence lattice of every subalgebra of $\mathbf{A}$.

If (δ, θ, ν) is a relevant triple of an algebra $\mathbf{B}$ such that (C1) holds in $\text{Con}(\mathbf{B})$, then $[\theta \wedge \nu, \nu] = [\theta, \nu] \leq \delta$ implies that $\theta \wedge [\nu, \nu] \leq \delta$, and since $\theta > \delta$ and δ is completely meet irreducible, it must be the case that $[\nu, \nu] \leq \delta$. Conversely, it is not hard to show (see the first paragraph of the proof of [10, Theorem 10.14]) that if (C1) fails in $\text{Con}(\mathbf{B})$, then there is a failure (with $x = \alpha$, $y = \beta$) of the following form:

$$[\alpha, \beta] \leq \delta < \theta \leq \alpha \leq [\beta, \beta] (\leq \beta)$$

where δ is completely meet irreducible and θ is its upper cover. It follows that $[\theta, \theta] \leq [\alpha, \beta] \leq \delta$, so θ/δ is abelian. Moreover, $\nu := (\delta : \theta) \geq \beta$ since $[\theta, \beta] \leq [\alpha, \beta] \leq \delta$, which implies that $[\nu, \nu] \geq [\beta, \beta] \geq \theta$. Consequently, (δ, θ, ν) is a relevant triple such that $[\nu, \nu] \not\leq \delta$. This shows that conditions (a)–(c) in Theorem 2.1 are also equivalent to the condition

- (d) $[\nu, \nu] \leq \delta$ for every relevant triple (δ, θ, ν) of a subalgebra of $\mathbf{A}$.

Corollary 2.2. *Let $\mathbf{A}$ be a finite algebra that generates a congruence modular variety $\mathcal{V}(\mathbf{A})$. If $\mathbf{A}$ satisfies the split centralizer condition, then (C1) holds in the congruence lattice of every subalgebra of $\mathbf{A}$, so $\mathcal{V}(\mathbf{A})$ is residually small.*

Proof. If $\mathbf{A}$ satisfies the split centralizer condition, and (δ, θ, ν) is a relevant triple of a subalgebra $\mathbf{B}$ of $\mathbf{A}$ which is split by the triple (α, β, κ) , then $[\![\delta, \nu]\!] \subseteq [\![\beta, \nu]\!] \searrow [\![\kappa, \alpha]\!]$ with the last interval abelian, therefore the other two intervals are also abelian. This proves that condition (d) holds for $\mathbf{A}$. Our statement now follows from the equivalence of (d) to conditions (c) and (b) in Theorem 2.1. $\square$

2.2. Compatible Relations, Entailment, and Dualizability. Let A be a set. By a *relation* on A we mean a subset ρ of A^n for some positive integer n , which we call the *arity* of ρ . For any nonempty subset I of $[n]$, *projection onto the coordinates in I* is the map $\text{proj}_I: A^n \rightarrow A^I$, $(a_i)_{i \in [n]} \mapsto (a_i)_{i \in I}$.

For any algebra $\mathbf{A}$, an n -ary relation ρ on A is called a *compatible relation of $\mathbf{A}$* if ρ is the universe of a subalgebra of $\mathbf{A}^n$ (or equivalently, ρ is a nonempty subuniverse of $\mathbf{A}^n$). The set of all compatible relations of $\mathbf{A}$ of arity $\leq n$ will be denoted by $\mathcal{R}_n(\mathbf{A})$, and $\mathcal{R}(\mathbf{A})$ will stand for the set $\bigcup_{n>0} \mathcal{R}_n(\mathbf{A})$ of all compatible relations of $\mathbf{A}$.

It is straightforward to check that if R is a set of compatible relations of an algebra $\mathbf{A}$, then so is every relation that can be obtained, in finitely many steps, from relations in $R \cup \{=\}$ by the following constructs:

- nonempty intersection of relations of the same arity,
- direct product of two relations,
- permutation of coordinates of a relation, and
- projection of a relation onto a nonempty subset of its coordinates.

The relations that can be obtained in this way are exactly the nonempty relations that are definable by primitive positive formulas (with $=$) using the relations in R . The relations that can be obtained, in finitely many steps, from relations in $R \cup \{=\}$ by the first three types of constructs are exactly the nonempty relations that are definable by quantifier free primitive positive formulas (with $=$) using the relations in R .

A *critical relation* of an algebra $\mathbf{A}$ is a compatible relation ρ of $\mathbf{A}$ that is

- (i) completely $\cap$ -irreducible in the lattice of subuniverses of $\mathbf{A}^n$ where n is the arity of ρ , and
- (ii) directly indecomposable as a relation, that is, $[n]$ cannot be partitioned into two nonempty sets I and J such that ρ and $\text{proj}_I(\rho) \times \text{proj}_J(\rho)$ differ only by a permutation of coordinates.

More informally, a compatible relation of $\mathbf{A}$ is critical if it cannot be obtained in a nontrivial way from other compatible relations using only the first three of the four types of constructs above.

In the theory of natural dualities there is an entailment concept, which we will denote by $\models_d$ (d stands for ‘duality’). We refer the reader interested in the definition to Definition 4.1 in [2, Chapter 2]. For the purposes of this paper an algebraic characterization of $\models_d$, which we state in Theorem 2.3 below, will be sufficient. We need some definitions beforehand.

Let B be a compatible relation of an algebra $\mathbf{A}$, say B is n -ary, and let $B' := \text{proj}_I(B)$ for some nonempty $I \subseteq [n]$. Then the projection map $\text{proj}_I: \mathbf{B} \rightarrow \mathbf{B}'$ is a surjective homomorphism between the algebras $\mathbf{B} (\leq \mathbf{A}^n)$ and $\mathbf{B}' (\leq \mathbf{A}^I)$ with universes B and B' . Following [2], we call $\text{proj}_I: \mathbf{B} \rightarrow \mathbf{B}'$ a *retractive projection* if it is a retraction, that is, if there exists a homomorphism $\varphi: \mathbf{B}' \rightarrow \mathbf{B}$ such that $\text{proj}_I \circ \varphi = \text{id}_{B'}$. An important special case is a *bijective projection* $\text{proj}_I: \mathbf{B} \rightarrow \mathbf{B}'$, when the retractive projection is bijective. Accordingly, we say that B' is obtained from B by retractive projection (or bijective projection) onto its coordinates in I .

From now on we will restrict to finite algebras. The following theorem is a consequence of Theorems 2.2 and 2.6 in [2, Chapter 9].

Theorem 2.3 (From [2]). *Let $\mathbf{A}$ be a finite algebra, and let $R \subseteq \mathcal{R}(\mathbf{A})$. For a nonempty relation ρ on A , we have $R \models_d \rho$ if and only if ρ can be obtained, in finitely many steps, from relations in $R \cup \{=\}$ by the following constructs:*

- *nonempty intersection of relations of the same arity,*
- *product of two relations,*
- *permutation of coordinates of a relation, and*
- *retractive projection of a relation onto a nonempty subset of its coordinates.*

The four types of constructs in this theorem will be referred to as $\models_d$ -constructs. Notice that the only difference between the list of $\models_d$ -constructs and the earlier list of

constructs for compatible relations is that among the $\models_d$ -constructs projections are restricted to retractive projections. Two immediate consequences are worth mentioning: (i) Since every $\models_d$ -construct occurs on the earlier list, it follows that if $R \subseteq \mathcal{R}(\mathbf{A})$ and $R \models_d \rho$, then $\rho \in \mathcal{R}(\mathbf{A})$. (ii) Since the first three constructs on the two lists are the same, our earlier remark implies that if $R \subseteq \mathcal{R}(\mathbf{A})$ and ρ is a nonempty relation on A that is definable by a quantifier-free primitive positive formula (with $=$) using relations in R , then $R \models_d \rho$.

Observation (i) above allows us to extend $\models_d$ to subsets of $\mathcal{R}(\mathbf{A})$ as follows: for $R, R' \subseteq \mathcal{R}(\mathbf{A})$ let $R \models_d R'$ mean that $R \models_d \rho$ for every $\rho \in R'$. It is easy to check that the relation $\models_d$ is transitive on subsets of $\mathcal{R}(\mathbf{A})$, that is, if $R, R', R'' \subseteq \mathcal{R}(\mathbf{A})$ satisfy $R \models_d R'$ and $R' \models_d R''$, then $R \models_d R''$.

The most powerful general criterion for dualizability is the following theorem of Willard and Zádori.

Theorem 2.4 (See [19],[20]). *Let $\mathbf{A}$ be a finite algebra. If $\mathcal{R}_n(\mathbf{A}) \models_d \mathcal{R}(\mathbf{A})$ for some integer $n \geq 1$, then $\mathbf{A}$ is dualizable.*

In fact, it is shown in [20] that if $\mathcal{R}_n(\mathbf{A}) \models_d \mathcal{R}(\mathbf{A})$ holds for some $n \geq 1$, then it also holds with the restriction that retractive projections among the $\models_d$ -constructs are limited to bijective projections.

2.3. Algebras with Parallelogram Terms. Let m and n be positive integers and let $k = m + n$. The concept of an (m, n) -parallelogram term (or k -parallelogram term) for a variety $\mathcal{V}$ was introduced in [13] to mean a term $\mathbf{P}$ such that the identities represented by the rows of the following matrix equation hold in $\mathcal{V}$:

$$(2.1) \quad \mathbf{P} \left(\begin{array}{ccc|cccccc} x & x & y & z & y & \cdots & y & y & \cdots & y & y \\ x & x & y & y & z & & y & y & & y & y \\ \vdots & & & \vdots & & \ddots & & & & \vdots & \\ x & x & y & y & y & & z & y & & y & y \\ y & x & x & y & y & & y & z & & y & y \\ \vdots & & & \vdots & & & & & \ddots & \vdots & \\ y & x & x & y & y & & y & y & & z & y \\ y & x & x & y & y & \cdots & y & y & \cdots & y & z \end{array} \right) = \begin{pmatrix} y \\ y \\ \vdots \\ y \\ y \\ \vdots \\ y \\ y \end{pmatrix}.$$

Here $\mathbf{P}$ is $(k+3)$ -ary, the rightmost block of variables is a $k \times k$ array, the upper left block is $m \times 3$ and the lower left block is $n \times 3$. An (m, n) -parallelogram term (or k -parallelogram term) for an algebra $\mathbf{A}$ is defined to be an (m, n) -parallelogram term (or k -parallelogram term) for the variety $\mathcal{V}(\mathbf{A})$ it generates.

It is easy to see from these definitions that a k -parallelogram term that is independent of its last k variables is a Maltsev term, and a k -parallelogram term that is independent of its first 3 variables is a k -ary near unanimity term.

It was proved in [13, Theorem 3.5] that if m, n, m', n' are positive integers such that $m + n = k = m' + n'$, then a variety has an (m, n) -parallelogram term if and only if it has an (m', n') -parallelogram term; this justifies referring to them as k -parallelogram terms. In addition, [13, Theorem 3.5] also shows that a variety has a k -parallelogram term if and only if it has a term, called a k -cube term, introduced in [1, Definition 2.4]. It follows from [1, Theorem 2.7] that every variety with a k -cube term is congruence modular. So, by combining all these results, or by a direct argument using [9, Theorem 3.2] we get the following conclusion.

Theorem 2.5 (See [1],[13];[9]). *Every variety with a parallelogram term is congruence modular.*

Our starting point for the proof of Theorem 1.1 will be the structure theorem in [13] (see [13, Theorems 4.1 and 2.5]) on the critical relations of a finite algebra with a parallelogram term. If C is an n -ary critical relation of $\mathbf{A}$, let $\mathbf{C}$ denote the subalgebra of $\mathbf{A}^n$ with universe C , and let $\mathbf{A}_i := \text{proj}_i(\mathbf{C})$ for each $i \in [n]$. Furthermore, let $\delta_i \in \text{Con}(\mathbf{A}_i)$ ($i \in [n]$) be such that $\delta := \prod_{i=1}^n \delta_i$ is the largest product congruence of $\prod_{i=1}^n \mathbf{A}_i$ with the property that $\mathbf{C}$ is a δ -saturated subalgebra of $\prod_{i=1}^n \mathbf{A}_i$. Then $\overline{\mathbf{C}} := \mathbf{C}/\delta_{\mathbf{C}}$ is a subdirect product of the algebras $\overline{\mathbf{A}}_i := \mathbf{A}_i/\delta_i$ ($i \in [n]$), which we call *the reduced representation of $\mathbf{C}$* .

The next theorem contains those parts of the structure theorem on critical relations that we will need later on; we retain the numbering from [13, Theorem 2.5].

Theorem 2.6 (From [13]). *Let C be an n -ary critical relation of a finite algebra $\mathbf{A}$ with a k -parallelogram term, and let $\overline{\mathbf{C}}$ be its reduced representation. If $n \geq k$ (> 1), then the following hold.*

- (1) $\overline{\mathbf{C}} \leq \prod_{i=1}^n \overline{\mathbf{A}}_i$ is a representation of $\overline{\mathbf{C}}$ as a subdirect product of subdirectly irreducible algebras $\overline{\mathbf{A}}_i$.
- (7) If $n > 2$, then each $\overline{\mathbf{A}}_i$ has abelian monolith μ_i ($i \in [n]$).
- (8) For the centralizers $\rho_\ell := (0 : \mu_\ell)$ of the monoliths μ_ℓ ($\ell \in [n]$), the image of the composite map

$$\overline{\mathbf{C}} \hookrightarrow \prod_{\ell=1}^n \overline{\mathbf{A}}_\ell \twoheadrightarrow \overline{\mathbf{A}}_i/\rho_i \times \overline{\mathbf{A}}_j/\rho_j$$

is the graph of an isomorphism for any $i, j \in [n]$.

Since $\overline{\mathbf{A}}_i = \mathbf{A}_i/\delta_i$ ($i \in [n]$) is subdirectly irreducible, $\delta_i \in \text{Con}(\mathbf{A}_i)$ is completely meet irreducible, and the monolith of $\overline{\mathbf{A}}_i$ is $\mu_i = \theta_i/\delta_i$ where $\theta_i \in \text{Con}(\mathbf{A}_i)$ is the unique cover of δ_i . Furthermore, $(0 : \mu_i) = \rho_i = \nu_i/\delta_i$ where $\nu_i = (\delta_i : \theta_i)$, and hence $\overline{\mathbf{A}}_i/\rho_i \cong \mathbf{A}_i/\nu_i$. Thus, with our current terminology of a relevant triple, Theorem 2.6 (restricted to the case $n > 2$) can be restated as follows.

Corollary 2.7. *Let C be an n -ary critical relation of a finite algebra $\mathbf{A}$ with a k -parallelogram term, and let $\mathbf{C}$ be the subalgebra of $\mathbf{A}^n$ with universe C . If $n \geq \max(3, k)$, then the following hold.*

- $\mathbf{C}$ is a subdirect product of the subalgebras $\mathbf{A}_i := \text{proj}_i(\mathbf{C})$ ($i \in [n]$) of $\mathbf{A}$.
- If $\delta = \prod_{i=1}^n \delta_i$ ($\delta_i \in \text{Con}(\mathbf{A}_i)$) is the largest product congruence of $\prod_{i=1}^n \mathbf{A}_i$ for which $\mathbf{C}$ is δ -saturated, then each δ_i ($i \in [n]$) is the first component of a relevant triple $(\delta_i, \theta_i, \nu_i)$ of $\mathbf{A}_i$.
- The assignment

$$\iota_{ij}: \mathbf{A}_i/\nu_i \rightarrow \mathbf{A}_j/\nu_j, \quad c_i/\nu_i \mapsto c_j/\nu_j \quad \text{whenever } (c_1, \dots, c_n) \in C$$

is an isomorphism for any $i, j \in [n]$.

3. REDUCTION TO ABELIAN CONGRUENCES OF BOUNDED INDEX

Our main tool for proving Theorem 1.1 will be Theorem 2.4. If $\mathbf{A}$ is a finite algebra satisfying the assumptions of Theorem 1.1, then Corollary 2.2 and Theorem 2.5 imply that (C1) holds in the congruence lattice of every subalgebra of $\mathbf{A}$. Hence ν'/δ' is abelian for every relevant triple (δ', θ', ν') of a subalgebra $\mathbf{A}'$ of $\mathbf{A}$. Therefore, if C is an n -ary critical relation of $\mathbf{A}$ such that, with the notation of Corollary 2.7, $\delta_i = 0$ for every $i \in [n]$, then ν_i is abelian for every $i \in [n]$, and hence the product congruence $\nu := \prod_{i=1}^n \nu_i$ of $\prod_{i=1}^n \mathbf{A}_i$ restricts to $\mathbf{C}$ as an abelian congruence $\nu_{\mathbf{C}}$. Moreover, by the last item in Corollary 2.7, $\nu_{\mathbf{C}}$ has index $< |A|$.

In this situation we can use modules associated to abelian congruences, as explained in Sections 4–5, to show that C is entailed by compatible relations of bounded arity. However, this argument does not work if, instead of an abelian interval $\llbracket 0, \nu_{\mathbf{C}} \rrbracket$, we have an abelian interval $\llbracket \delta_{\mathbf{C}}, \nu_{\mathbf{C}} \rrbracket$ with $\delta_{\mathbf{C}} \neq 0$. In the general case when $\delta_{\mathbf{C}}$ may be a nontrivial congruence of $\mathbf{C}$, we will use the assumption that $\mathbf{A}$ satisfies the split centralizer condition to replace C by another compatible relation B , which is not weaker than C with respect to entailment, but has an abelian interval $\llbracket 0, \tilde{\alpha}_{\mathbf{C}} \rrbracket$ corresponding to $\llbracket \delta_{\mathbf{C}}, \nu_{\mathbf{C}} \rrbracket$ at the bottom of the congruence lattice, moreover, the index of $\tilde{\alpha}_{\mathbf{C}}$ remains bounded by a number independent of n (though might be much bigger than the index of $\nu_{\mathbf{C}}$, which is $< |A|$).

The purpose of this section is to construct such a relation B for every critical relation C of $\mathbf{A}$.

Given a finite algebra $\mathbf{A}$, we define several constants associated to $\mathbf{A}$ as follows.

- Let $\mathbf{a}$ be the maximum of all $|\text{Aut}(\mathbf{S}/\nu)|$ where $\mathbf{S} \leq \mathbf{A}$ and (δ, θ, ν) is a relevant triple of $\mathbf{S}$. ($\mathbf{a}$ stands for ‘automorphisms’.)
- Let $\mathbf{s}$ be the number of distinct pairs $(\mathbf{S}, \delta)$ such that $\mathbf{S} \leq \mathbf{A}$ and (δ, θ, ν) is a relevant triple of $\mathbf{S}$. ($\mathbf{s}$ stands for ‘subdirectly irreducible sections $\mathbf{S}/\delta$ ’.)
- Let $\mathbf{i} := |A|^{\mathbf{as}}$. ($\mathbf{i}$ stands for ‘index’; see Theorem 3.1 below.)

- Let $\mathbf{p}$ be the least positive integer with the property that for every subalgebra $\mathbf{S} \leq \mathbf{A}$ every relevant triple of $\mathbf{S}$ is split by a triple (α, β, κ) such that $\mathbf{S}/\kappa$ embeds into $\mathbf{A}^{\mathbf{p}}$ for some $p \leq \mathbf{p}$. ($\mathbf{p}$ stands for ‘power’.)

Theorem 3.1. *Let $\mathbf{A}$ be a finite algebra with a k -parallelogram term such that $\mathbf{A}$ satisfies the split centralizer condition. For every $n \geq \max(3, k)$ and for every n -ary critical relation C of $\mathbf{A}$, there exists a compatible relation B of $\mathbf{A}$ which has the following two properties:*

- (*) *There exist*
 - (I) *subalgebras $\mathbf{B}_i \leq \mathbf{A}^{p_i}$ with $p_i \leq \mathbf{p}$ for each $i \in [n]$ such that $\mathbf{B}_i$ is isomorphic to a section of $\mathbf{A}$, and*
 - (II) *nontrivial abelian congruences $\tilde{\alpha}_i \in \text{Con}(\mathbf{B}_i)$ ($i \in [n]$) such that*
 - (III) *B is the universe of a subdirect product $\mathbf{B}$ of $\mathbf{B}_1, \dots, \mathbf{B}_n$, and*
 - (IV) *the product congruence $\tilde{\alpha} := \prod_{i=1}^n \tilde{\alpha}_i$ of $\prod_{i=1}^n \mathbf{B}_i$ restricts to $\mathbf{B}$ as a congruence $\tilde{\alpha}_{\mathbf{B}}$ of index $\leq \mathbf{i}$.*
- (**) $\mathcal{R}_{1+\mathbf{p}}(\mathbf{A}) \cup \{B\} \models_d C$.

Proof. Corollary 2.7 describes the structure of C . Using the notation of Corollary 2.7, we first define two equivalence relations, $\approx$ and $\sim$, on $[n]$ as follows:

- $i \approx j$ iff $\mathbf{A}_i = \mathbf{A}_j$ and $(\delta_i, \theta_i, \nu_i) = (\delta_j, \theta_j, \nu_j)$;
- $i \sim j$ iff $i \approx j$ and ι_{ij} is the identity isomorphism.

To give an upper bound for the indices of $\approx$ and $\sim$, notice that if $i, j \in [n]$ are such that $\mathbf{A}_i = \mathbf{A}_j$ and $\delta_i = \delta_j$, then it follows that $(\delta_i, \theta_i, \nu_i) = (\delta_j, \theta_j, \nu_j)$, so $i \approx j$. This implies that $\approx$ has at most $\mathbf{s}$ classes, that is, $\approx$ has index $|[n]/\approx| \leq \mathbf{s}$. The definition of $\sim$ shows that every class of $\approx/\sim$ has size $\leq \mathbf{a}$. Hence $\sim$ has at most $\mathbf{as}$ classes, that is, $\sim$ has index $|[n]/\sim| \leq \mathbf{as}$.

Our assumption that $\mathbf{A}$ satisfies the split centralizer condition ensures that for each $i \in [n]$ the relevant triple $(\delta_i, \theta_i, \nu_i)$ of $\mathbf{A}_i$ ($\leq \mathbf{A}$) is split by a triple $(\alpha_i, \beta_i, \kappa_i)$. These choices could differ coordinate by coordinate, but we can choose a transversal $T_{\approx}$ for the classes of $\approx$ and redefine the triples in coordinates $i \notin T_{\approx}$ to arrange that $(\alpha_i, \beta_i, \kappa_i) = (\alpha_j, \beta_j, \kappa_j)$ whenever $i \approx j$ ($i, j \in [n]$).

Let α, β, δ , and κ denote the product congruences $\prod_{i=1}^n \alpha_i, \prod_{i=1}^n \beta_i, \prod_{i=1}^n \delta_i$, and $\prod_{i=1}^n \kappa_i$ of $\prod_{i=1}^n \mathbf{A}_i$. We have $\kappa \leq \alpha, \kappa \leq \beta \leq \delta$, and $\kappa = \alpha \wedge \beta$, because $\kappa_i \leq \alpha_i, \kappa_i \leq \beta_i \leq \delta_i$, and $\kappa_i = \alpha_i \wedge \beta_i$ for all i . By Corollary 2.7, $\mathbf{C}$ is δ -saturated, and hence also β - and κ -saturated.

Now let D denote the set of all tuples $(d_1, \dots, d_n) \in C$ such that

- (†) $d_i \equiv_{\alpha_j} d_j$ whenever $i \sim j$.

Note that $i \sim j$ implies that $\alpha_i = \alpha_j$, so α_i and α_j are interchangeable in (†).

Claim 3.2. *Let T be a transversal for the classes of $\sim$. For every $(c_1, \dots, c_n) \in C$*

- (1) *there exists $(d_1, \dots, d_n) \in D$ such that*
 $(\dagger) \ (d_1, \dots, d_n) \equiv_\beta (c_1, \dots, c_n)$ *and $d_t = c_t$ for all $t \in T$;*
- (2) *the tuples $(d_1, \dots, d_n) \in D$ satisfying $(\dagger)$ are uniquely determined modulo κ .*

Proof of Claim 3.2. Choose $(c_1, \dots, c_n) \in C$.

For (1), define $d_t := c_t$ for $t \in T$. Our aim is to show that

- (3.1) for each $t \in T$ and for all $i \sim t$ with $i \neq t$
there exist $d_i \in A_i$ such that $c_i \equiv_{\beta_i} d_i \equiv_{\alpha_t} d_t$.

This will complete the proof of (1) for the following reason. Let $(d_1, \dots, d_n)$ be a tuple obtained in this way. By its construction, $(d_1, \dots, d_n)$ will satisfy condition $(\dagger)$. It will also satisfy condition $(\dagger)$, because whenever $i \sim j$, we have a $t \in T$ with $i, j \sim t$, so $d_i, d_j \equiv_{\alpha_t} d_t$; since $\alpha_j = \alpha_t$, we get that $d_i \equiv_{\alpha_j} d_j$. Now, since C is β -saturated, the fact that $(\dagger)$ holds for $(d_1, \dots, d_n)$ ensures that $(d_1, \dots, d_n) \in C$. Hence the fact that $(\dagger)$ also holds for $(d_1, \dots, d_n)$ implies that $(d_1, \dots, d_n) \in D$.

To prove (3.1), choose $t \in T$ and $i \sim t$ with $i \neq t$. Then $i \sim t$ implies that $\mathbf{A}_i = \mathbf{A}_t$, $\beta_i = \beta_t$, $\nu_i = \nu_t$, and $c_i/\nu_i = c_t/\nu_t$. Hence, $c_i \equiv_{\nu_t} c_t$. By our assumptions, $(\alpha_t, \beta_t, \kappa_t)$ splits $(\delta_t, \theta_t, \nu_t)$, so $\alpha_t \vee \beta_t = \nu_t$ and α_t/κ_t is an abelian congruence of $\mathbf{A}_t/\kappa_t$. Since $\mathbf{A}_t/\kappa_t$ lies in a congruence modular variety, we have that α_t/κ_t permutes with all congruences of $\mathbf{A}_t/\kappa_t$ (see [10, Theorem 6.2]). Hence, $\nu_t = \alpha_t \vee \beta_t = \beta_t \circ \alpha_t$. Thus, the fact that $c_i \equiv_{\nu_t} c_t$ holds implies that there exists $d_i \in A_t = A_i$ such that $c_i \equiv_{\beta_t} d_i \equiv_{\alpha_t} c_t = d_t$. Since $\beta_i = \beta_t$, this d_i is the element we wanted to find.

For (2), assume that $(d_1, \dots, d_n), (d'_1, \dots, d'_n) \in D$ both satisfy condition $(\dagger)$. We want to show that $(d_1, \dots, d_n) \equiv_\kappa (d'_1, \dots, d'_n)$. Since $\kappa = \alpha \wedge \beta$, this is equivalent to showing that $(d_1, \dots, d_n) \equiv_\alpha (d'_1, \dots, d'_n)$ and $(d_1, \dots, d_n) \equiv_\beta (d'_1, \dots, d'_n)$. The latter follows from the assumption that both tuples satisfy condition $(\dagger)$. To prove the former, consider any $i \in [n]$, and let $t \in T$ be such that $i \sim t$. Combining the second part of condition $(\dagger)$ with condition $(\dagger)$ in the definition of D , we get that $c_t = d_t \equiv_{\alpha_i} d_i$, and similarly, $c_t = d'_t \equiv_{\alpha_i} d'_i$, hence $d_i \equiv_{\alpha_i} d'_i$. Thus, $(d_1, \dots, d_n) \equiv_\alpha (d'_1, \dots, d'_n)$, completing the proof of (2). $\diamond$

Claim 3.3. *D is the universe of a subalgebra $\mathbf{D}$ of $\mathbf{C}$ with the following properties:*

- (1) $\mathbf{D}$ is a subdirect product of $\mathbf{A}_1, \dots, \mathbf{A}_n$,
- (2) $\text{proj}_T(\mathbf{D}) = \text{proj}_T(\mathbf{C})$ for every transversal T for the classes of $\sim$,
- (3) $\mathbf{D}[\beta] = \mathbf{C}$, and
- (4) $\mathbf{D}[\kappa] = \mathbf{D}$.

Proof of Claim 3.3. Consider a transversal T for the classes of $\sim$. Then $D \subseteq C$ implies that $\text{proj}_T(D) \subseteq \text{proj}_T(C)$, and Claim 3.2 (1) shows that $\text{proj}_T(D) \supseteq \text{proj}_T(C)$. Thus, $\text{proj}_T(D) = \text{proj}_T(C)$. This implies that $\text{proj}_t(D) = \text{proj}_t(C) = A_t$ for every $t \in T$. Since every element of $[n]$ occurs in some transversal T , we get that $\text{proj}_i(D) = A_i$ for all $i \in [n]$. In particular, we see that $D \neq \emptyset$. The definition of D shows that

$D \subseteq C$ and D is definable by a primitive positive formula, using the compatible relations $\alpha_1, \dots, \alpha_n$ and C of $\mathbf{A}$. Hence, D is a compatible relation of $\mathbf{A}$. These arguments prove that D is the universe of a subalgebra $\mathbf{D}$ of $\mathbf{C}$ satisfying (1)–(2).

For (3), observe that since $\mathbf{D} \leq \mathbf{C}$ and $\mathbf{C}$ is β -saturated, it follows that $\mathbf{D}[\beta] \leq \mathbf{C}[\beta]$ and $\mathbf{C}[\beta] = \mathbf{C}$. Thus, $\mathbf{D}[\beta] \leq \mathbf{C}$. However, we get from Claim 3.2 (1) that $C \subseteq D[\beta]$. Hence, $\mathbf{D}[\beta] = \mathbf{C}$, as claimed.

Finally, we prove (4). Clearly, $\mathbf{D} \leq \mathbf{D}[\kappa]$. The reverse inclusion $D[\kappa] \subseteq D$ can be established as follows:

$$D[\kappa] \subseteq D[\alpha] \cap D[\beta] = D[\alpha] \cap C \subseteq D,$$

where the last inclusion is an immediate consequence of the definition of D . This completes the proof of Claim 3.3. $\diamond$

Claim 3.4. *For every transversal T for the classes of $\sim$,*

$$|\mathbf{D}/\alpha_{\mathbf{D}}| \leq |\text{proj}_T(\mathbf{D})| \leq i.$$

Proof of Claim 3.4. Let T be a transversal for the classes of $\sim$. From the inequality $|[n]/\sim| \leq \text{as}$ proved earlier we get that $|T| \leq \text{as}$. Thus, $|\text{proj}_T(\mathbf{D})| \leq |A|^{|T|} \leq |A|^{\text{as}} = i$, which establishes the second inequality.

It follows from the definition of D that the kernel of the projection map $\text{proj}_T: \mathbf{D} \rightarrow \text{proj}_T(\mathbf{D})$ is contained in α . This implies the first inequality in Claim 3.4. $\diamond$

Now we are ready to define the algebras $\mathbf{B}_1, \dots, \mathbf{B}_n$, $\mathbf{B}$, and the congruences $\tilde{\alpha}_i \in \text{Con}(\mathbf{B}_i)$ for which the conclusions (*)–(**) of Theorem 3.1 hold.

Since each κ_i is a $\mathcal{Q}$ -congruence of $\mathbf{A}_i$ for $\mathcal{Q} = \text{SP}(\mathbf{A})$, we have that there exist subalgebras $\mathbf{B}_i \leq \mathbf{A}^{p_i}$ with $p_i \leq \mathbf{p}$ for each i , and surjective homomorphisms $\varphi_i: \mathbf{A}_i \rightarrow \mathbf{B}_i$ with kernels κ_i ($i \in [n]$) such that φ_i induces an isomorphism $\bar{\varphi}_i: \mathbf{A}_i/\kappa_i \rightarrow \mathbf{B}_i$. Recall that we chose the splitting triples so that $(\alpha_i, \beta_i, \kappa_i) = (\alpha_j, \beta_j, \kappa_j)$ whenever $i \approx j$ (and hence $\mathbf{A}_i = \mathbf{A}_j$). With the same reasoning, we can arrange that the algebras $\mathbf{B}_i$ and the homomorphisms φ_i are selected so that $\mathbf{B}_i = \mathbf{B}_j$ and $\varphi_i = \varphi_j$ whenever $i \approx j$. For every $i \in [n]$, define $\tilde{\alpha}_i \in \text{Con}(\mathbf{B}_i)$ to be the image of $\alpha_i \in \text{Con}(\mathbf{A}_i)$ under the homomorphism φ_i , which is the same as the image of $\alpha_i/\kappa_i \in \text{Con}(\mathbf{A}_i/\kappa_i)$ under the isomorphism $\bar{\varphi}_i$; so $\tilde{\alpha}_i$ is indeed a congruence of $\mathbf{B}_i$. Furthermore, define $\mathbf{B}$ to be the image of $\mathbf{D}$ under the product homomorphism $\varphi := \prod_{i=1}^n \varphi_i: \prod_{i=1}^n \mathbf{A}_i \rightarrow \prod_{i=1}^n \mathbf{B}_i$.

Claim 3.5. *Conditions (I)–(IV) hold for the algebras $\mathbf{B}_1, \dots, \mathbf{B}_n$ and $\mathbf{B}$, and for the congruences $\tilde{\alpha}_1, \dots, \tilde{\alpha}_n$ defined above. Moreover, $\mathbf{D}$ is the full inverse image of $\mathbf{B}$ under the homomorphism φ .*

Proof of Claim 3.5. (I) follows from the choice of the $\mathbf{B}_i$'s.

(II) holds, because for every $i \in [n]$, $\tilde{\alpha}_i = \bar{\varphi}_i[\alpha_i/\kappa_i]$ where $\bar{\varphi}_i: \mathbf{A}_i/\kappa_i \rightarrow \mathbf{B}_i$ is an isomorphism and α_i/κ_i is a nontrivial abelian congruence of $\mathbf{A}_i$. (The latter follows from the fact that $(\alpha_i, \beta_i, \kappa_i)$ splits the relevant triple $(\delta_i, \theta_i, \nu_i)$ of $\mathbf{A}_i$.)

For (III), recall that $\mathbf{D}$ is a subdirect product of $\mathbf{A}_1, \dots, \mathbf{A}_n$ by Claim 3.3(1), and the homomorphisms $\varphi_i: \mathbf{A}_i \rightarrow \mathbf{B}_i$ ($i \in [n]$) are onto. Since $\varphi = \prod_{i=1}^n \varphi_i$, it follows that $\mathbf{B} = \varphi[\mathbf{D}]$ is a subdirect product of $\mathbf{B}_1, \dots, \mathbf{B}_n$.

To verify (IV) notice that $\varphi := \prod_{i=1}^n \varphi_i: \prod_{i=1}^n \mathbf{A}_i \rightarrow \prod_{i=1}^n \mathbf{B}_i$ is a surjective homomorphism with kernel $\kappa = \prod_{i=1}^n \kappa_i$, therefore φ decomposes as shown by the first line of the array below:

$$\begin{array}{ccccccc}
 \varphi: & \prod_{i=1}^n \mathbf{A}_i & \xrightarrow{\text{nat}} & (\prod_{i=1}^n \mathbf{A}_i)/\kappa & \xrightarrow{\cong} & \prod_{i=1}^n (\mathbf{A}_i/\kappa_i) & \xrightarrow{\bar{\varphi}} & \prod_{i=1}^n \mathbf{B}_i \\
 & \alpha & \mapsto & \alpha/\kappa & \mapsto & \prod_{i=1}^n \alpha_i/\kappa_i & \mapsto & \tilde{\alpha} \\
 (3.2) & & & & & & & \\
 \varphi|_{\mathbf{D}}: & \mathbf{D} & \xrightarrow{\text{nat}} & \mathbf{D}/\kappa_{\mathbf{D}} & \xrightarrow{\bar{\varphi} \circ \cong} & & & \mathbf{B} \\
 & \alpha_{\mathbf{D}} & \mapsto & \alpha_{\mathbf{D}}/\kappa_{\mathbf{D}} & \mapsto & & & \tilde{\alpha}_{\mathbf{B}}
 \end{array}$$

The leftmost factor of φ is the natural map, the middle factor is the natural isomorphism, and the rightmost factor is the isomorphism $\bar{\varphi} := \prod_{i=1}^n \bar{\varphi}_i$. Since $\tilde{\alpha}_i = \varphi[\alpha_i]$ for every $i \in [n]$, φ maps the congruence $\alpha = \prod_{i=1}^n \alpha_i$ of $\prod_{i=1}^n \mathbf{A}_i$ onto the congruence $\tilde{\alpha} = \prod_{i=1}^n \tilde{\alpha}_i$ of $\prod_{i=1}^n \mathbf{B}_i$, via this factorization, as indicated by the second line of (3.2). Therefore, combining the two isomorphisms among the factors of φ and restricting φ to $\mathbf{D}$ yields that $\varphi|_{\mathbf{D}}$ factors and acts on $\alpha_{\mathbf{D}}$ as the third and fourth lines of (3.2) show. Thus,

$$\mathbf{D}/\alpha_{\mathbf{D}} \cong (\mathbf{D}/\kappa_{\mathbf{D}})/(\alpha_{\mathbf{D}}/\kappa_{\mathbf{D}}) \cong \mathbf{B}/\tilde{\alpha}_{\mathbf{B}}.$$

This implies that the index of $\tilde{\alpha}_{\mathbf{B}}$ in $\mathbf{B}$ is $|\mathbf{B}/\tilde{\alpha}_{\mathbf{B}}| = |\mathbf{D}/\alpha_{\mathbf{D}}|$, which is $\leq i$ by Claim 3.4.

For the last statement of the claim, $B = \varphi[D]$ implies that $\varphi^{-1}[B] \supseteq D$. Since the kernel of φ is κ , the equality $B = \varphi[D]$ also implies that $\varphi^{-1}[B] \subseteq D[\kappa]$. But we know from Claim 3.3 that $D[\kappa] = D$, so the equality $\varphi^{-1}[B] = D$ we wanted to prove follows. $\diamond$

Claim 3.6. *For every $i \in [n]$, A_i , B_i , the (graph of the) homomorphism φ_i , and*

$$\beta_i \circ \varphi_i := \{(x, y) \in A_i \times B_i : x \equiv_{\beta_i} z \text{ and } y = \varphi_i(z) \text{ for some } z \in A_i\}$$

are compatible relations of $\mathbf{A}$. Moreover,

$$(3.3) \quad \{A_i, B_i, \varphi_i, \beta_i \circ \varphi_i : i \in [n]\} \cup \{B\} \models_d C.$$

Proof of Claim 3.6. Choose any $i \in [n]$. Since $\mathbf{A}_i \leq \mathbf{A}$ and $\mathbf{B}_i \leq \mathbf{A}^{p_i}$, we have that $A_i \in \mathcal{R}_1(\mathbf{A})$ and $B_i \in \mathcal{R}_{p_i}(\mathbf{A})$. The fact that φ_i is a homomorphism $\mathbf{A}_i \rightarrow \mathbf{B}_i$ implies that its graph is the universe of a subalgebra of $\mathbf{A}_i \times \mathbf{B}_i \leq \mathbf{A}^{1+p_i}$. Thus, $\varphi_i \in \mathcal{R}_{1+p_i}(\mathbf{A})$. The definition of $\beta_i \circ \varphi_i$ shows that $\beta_i \circ \varphi_i \neq \emptyset$ and $\beta_i \circ \varphi_i$ is definable by a primitive positive formula, using the compatible relations $A_i, B_i, \beta_i, \varphi_i \in \mathcal{R}(\mathbf{A})$, therefore $\beta_i \circ \varphi_i \in \mathcal{R}(\mathbf{A})$. In fact, $\beta_i \circ \varphi_i$ is also the universe of a subalgebra of $\mathbf{A}_i \times \mathbf{B}_i \leq \mathbf{A}^{1+p_i}$, so $\beta_i \circ \varphi_i \in \mathcal{R}_{1+p_i}(\mathbf{A})$. This proves the first statement of the claim.

To prove the second statement, let us fix a transversal T for the classes of $\sim$, and define X to be the set of all tuples

$$(c_1, \dots, c_n, b_1, \dots, b_n) \in \prod_{i=1}^n A_i \times \prod_{i=1}^n B_i \left(\subseteq A^{n+\sum p_i} \right)$$

such that

- $(b_1, \dots, b_n) \in B$,
- $(c_i, b_i) \in \beta_i \circ \varphi_i$ for all $i \in [n]$, and
- $b_t = \varphi_t(c_t)$ for all $t \in T$.

Our main goal is to show that

- (i) $\text{proj}_{[n]}(X) = C$ and
- (ii) the projection map $\text{proj}_{[n]}: X \rightarrow C$ is one-to-one,

because we can deduce (3.3) from statements (i)–(ii) as follows. We have $X \neq \emptyset$ by (i), and X is definable by a quantifier-free primitive positive formula using the relations A_i , B_i , B , φ_t , and $\beta_i \circ \varphi_i$, therefore

$$\{A_i, B_i, \varphi_i, \beta_i \circ \varphi_i : i \in [n]\} \cup \{B\} \models_d X.$$

Furthermore, by (i) and (ii), C can be obtained from X by bijective projection, which is an $\models_d$ -construct, so $\{X\} \models_d C$. Hence (3.3) follows by the transitivity of $\models_d$.

It remains to prove (i) and (ii). For the inclusion $\supseteq$ in (i), let $(c_1, \dots, c_n) \in C$. By Claim 3.2, there exists $(d_1, \dots, d_n) \in D$ such that $(d_1, \dots, d_n) \equiv_\beta (c_1, \dots, c_n)$ and $d_t = c_t$ for all $t \in T$. Let $b_i = \varphi_i(d_i)$ for each $i \in [n]$. By the definition of B , this choice ensures that $(b_1, \dots, b_n) \in B$. Furthermore, $c_i \equiv_{\beta_i} d_i$ and $b_i = \varphi_i(d_i)$ imply $(c_i, b_i) \in \beta_i \circ \varphi_i$ for every $i \in [n]$, while $d_t = c_t$ and $b_t = \varphi_t(d_t)$ imply $b_t = \varphi_t(c_t)$ for every $t \in T$. Hence, $(c_1, \dots, c_n, b_1, \dots, b_n) \in X$, so $(c_1, \dots, c_n) \in \text{proj}_{[n]}(X)$.

The inclusion $\subseteq$ in (i) and the claim in (ii) will follow if we prove the following statement:

for every $(c_1, \dots, c_n) \in \text{proj}_{[n]}(X)$ we have that

- $(c_1, \dots, c_n) \in C$, and
- there is a unique tuple $(b_1, \dots, b_n)$ such that $(c_1, \dots, c_n, b_1, \dots, b_n) \in X$.

So, let $(c_1, \dots, c_n) \in \text{proj}_{[n]}(X)$. Then $(c_1, \dots, c_n, b_1, \dots, b_n) \in X$ for at least one tuple $(b_1, \dots, b_n)$. By the definition of X it must be the case that $(b_1, \dots, b_n) \in B$, $(c_i, b_i) \in \beta_i \circ \varphi_i$ for all $i \in [n]$, and $b_t = \varphi_t(c_t)$ for all $t \in T$. For any $t \in T$, define $d_t := c_t$; so $c_t \equiv_{\beta_t} d_t$ and $b_t = \varphi_t(d_t)$. For $i \in [n] \setminus T$, use the definition of $\beta_i \circ \varphi_i$ to get d_i such that $c_i \equiv_{\beta_i} d_i$ and $b_i = \varphi_i(d_i)$. Thus, $(c_1, \dots, c_n) \equiv_\beta (d_1, \dots, d_n)$ and $(d_1, \dots, d_n) \in \varphi^{-1}[(b_1, \dots, b_n)]$. We established in Claim 3.5 that $D = \varphi^{-1}[B]$, so $(d_1, \dots, d_n) \in \varphi^{-1}[(b_1, \dots, b_n)]$ implies that $(d_1, \dots, d_n) \in D$. Since $D \subseteq C$ and C is β -saturated, $(c_1, \dots, c_n) \equiv_\beta (d_1, \dots, d_n)$ yields that $(c_1, \dots, c_n) \in C$.

For the uniqueness of $(b_1, \dots, b_n)$ observe that our argument in the preceding paragraph shows the following: if $(b_1, \dots, b_n)$ is such that $(c_1, \dots, c_n, b_1, \dots, b_n) \in X$, then

there exists $(d_1, \dots, d_n) \in D$ with the properties

$$(c_1, \dots, c_n) \equiv_\beta (d_1, \dots, d_n) \in \varphi^{-1}[(b_1, \dots, b_n)] \quad \text{and} \quad c_t = d_t \text{ for all } t \in T.$$

So, if $(b'_1, \dots, b'_n)$ is another tuple with $(c_1, \dots, c_n, b'_1, \dots, b'_n) \in X$, then there exists $(d'_1, \dots, d'_n) \in D$ such that

$$(c_1, \dots, c_n) \equiv_\beta (d'_1, \dots, d'_n) \in \varphi^{-1}[(b'_1, \dots, b'_n)] \quad \text{and} \quad c_t = d'_t \text{ for all } t \in T.$$

It follows from Claim 3.2(2) that $(d_1, \dots, d_n) \equiv_{\kappa_{\mathbf{D}}} (d'_1, \dots, d'_n)$. Since $\kappa_{\mathbf{D}}$ is the kernel of φ we get that

$$(b_1, \dots, b_n) = \varphi((d_1, \dots, d_n)) = \varphi((d'_1, \dots, d'_n)) = (b'_1, \dots, b'_n),$$

proving the uniqueness of $(b_1, \dots, b_n)$. $\diamond$

Statement (*) of Theorem 3.1 was proved in Claim 3.5. Statement (**) of Theorem 3.1 follows from Claim 3.6 and the fact (established in the proof of Claim 3.6) that each one of the relations $A_i, B_i, \varphi_i, \beta_i \circ \varphi_i$ ($i \in [n]$) is a member of $\mathcal{R}_m(\mathbf{A})$ for some $m \leq 1 + \mathbf{p}$.

This completes the proof of Theorem 3.1. $\square$

4. ABELIAN CONGRUENCES AND MODULES

In [10, Chapter 9] Freese and McKenzie describe how matrix rings of any size can be associated to a congruence modular variety $\mathcal{V}$, and then modules over these rings to any abelian congruence α of an algebra $\mathbf{C}$ in $\mathcal{V}$. They prove in [10, Theorem 9.9] that there is a strong connection between the pair $(\mathbf{C}, \alpha)$ and the associated modules, namely, if all α -classes are represented in the module, then the interval $\llbracket 0, \alpha \rrbracket$ in $\text{Con}(\mathbf{C})$ is isomorphic to the lattice of submodules of the associated module.

Our goal in this section is to prove an analogous result for subalgebras in place of congruences. We will start by recalling the relevant definitions and basic facts from [10, Chapter 9], but we will slightly change the notation.

Throughout this section we will work under the following global assumptions:

- $\mathcal{V}$ is a congruence modular variety,
- d is a fixed difference term for $\mathcal{V}$, and
- $\mathcal{O}$ is a fixed nonempty set of constant symbols not occurring in the language of $\mathcal{V}$.

Recall that every congruence modular variety has a difference term (see [10, Theorem 5.5]), which is a ternary term d satisfying the following conditions for every algebra $\mathbf{C} \in \mathcal{V}$ and congruence ϑ of $\mathbf{C}$:

$$(4.1) \quad d^{\mathbf{C}}(x, x, y) = y \quad \text{for all } x, y \in \mathbf{C}, \text{ and}$$

$$(4.2) \quad d^{\mathbf{C}}(y, x, x) \equiv_{[\vartheta, \vartheta]} y \quad \text{for all } x, y \in \mathbf{C} \text{ such that } x \equiv_{\vartheta} y.$$

In addition, d also satisfies the following condition (see [10, Proposition 5.7]): for every term $t = t(x_1, \dots, x_k)$ and for arbitrary abelian congruence α of an algebra $\mathbf{C} \in \mathcal{V}$,

$$(4.3) \quad \begin{aligned} d^{\mathbf{C}}(t^{\mathbf{C}}(u_1, \dots, u_k), t^{\mathbf{C}}(v_1, \dots, v_k), t^{\mathbf{C}}(w_1, \dots, w_k)) \\ = t^{\mathbf{C}}(d^{\mathbf{C}}(u_1, v_1, w_1), \dots, d^{\mathbf{C}}(u_k, v_k, w_k)) \end{aligned}$$

whenever $u_i, v_i, w_i \in \mathbf{C}$ are such that $u_i \equiv_{\alpha} v_i \equiv_{\alpha} w_i$ for all i .

We will use the set $\mathcal{O}$ of new constant symbols to expand the language of $\mathcal{V}$. Terms in the expanded language will be called $\mathcal{O}$ -terms, and the algebras obtained from the members of $\mathcal{V}$ by interpreting all constant symbols $o \in \mathcal{O}$ will be called $\mathcal{O}$ -algebras. The class of algebras obtained in this way will be denoted by $\mathcal{V}_{\mathcal{O}}$. We will restrict the use of the phrases ‘subalgebra’, ‘homomorphism’, and ‘product’ to the algebras in $\mathcal{V}$ (i.e., to algebras in the original language), and use the phrases ‘ $\mathcal{O}$ -subalgebra’, ‘ $\mathcal{O}$ -homomorphism’, and ‘ $\mathcal{O}$ -product’ for $\mathcal{O}$ -algebras. There will be no need for using the phrase ‘ $\mathcal{O}$ -congruence’, because every $\mathcal{O}$ -algebra has the same congruence lattice, with the same commutator operation, as its reduct to the language of $\mathcal{V}$.

If we apply Corollary 5.8 (along with Lemma 5.6) from [10] to $\mathcal{O}$ -algebras in $\mathcal{V}_{\mathcal{O}}$, we get the following.

Lemma 4.1 (From [10]). *If α is an abelian congruence of an $\mathcal{O}$ -algebra $\mathbf{C}$ in $\mathcal{V}_{\mathcal{O}}$, then for every $o \in \mathcal{O}$, the α -class containing $o^{\mathbf{C}}$ is an abelian group*

$$\mathbf{M}_{\mathbf{C}}(\alpha, o) := (o^{\mathbf{C}}/\alpha; +_o, -_o, o^{\mathbf{C}})$$

with zero element $o^{\mathbf{C}}$ for the operations $+_o$ and $-_o$ defined by

$$u +_o v := d^{\mathbf{C}}(u, o^{\mathbf{C}}, v) \quad \text{and} \quad -_o u := d^{\mathbf{C}}(o^{\mathbf{C}}, u, o^{\mathbf{C}}) \quad \text{for all } u, v \in o^{\mathbf{C}}/\alpha.$$

Furthermore,

- (i) $d^{\mathbf{C}}(u, v, w) = u -_o v +_o w$ for all $u, v, w \in o^{\mathbf{C}}/\alpha$, and
- (ii) the $\mathcal{O}$ -term operations of $\mathbf{C}$ are linear between the α -classes; more precisely, if $r(x_1, \dots, x_k)$ is an $\mathcal{O}$ -term and $o_1, \dots, o_k, o \in \mathcal{O}$ are such that $r(o_1, \dots, o_k) = o$, then

$$r^{\mathbf{C}}|_{(o_1^{\mathbf{C}}/\alpha) \times \dots \times (o_k^{\mathbf{C}}/\alpha)} : \mathbf{M}_{\mathbf{C}}(\alpha, o_1) \times \dots \times \mathbf{M}_{\mathbf{C}}(\alpha, o_k) \rightarrow \mathbf{M}_{\mathbf{C}}(\alpha, o)$$

is a group homomorphism. Consequently, for the unary $\mathcal{O}$ -terms $r_i(x) := r(o_1, \dots, o_{i-1}, x, o_{i+1}, \dots, o_k)$ ($1 \leq i \leq k$) the maps $r_i^{\mathbf{C}}|_{o_i^{\mathbf{C}}/\alpha} : \mathbf{M}_{\mathbf{C}}(\alpha, o_i) \rightarrow \mathbf{M}_{\mathbf{C}}(\alpha, o)$ are also group homomorphisms, and

$$(4.4) \quad \begin{aligned} r^{\mathbf{C}}(u_1, \dots, u_k) &= r_1^{\mathbf{C}}(u_1) +_o \dots +_o r_k^{\mathbf{C}}(u_k) \\ &\text{for all } u_i \in o_i^{\mathbf{C}}/\alpha \text{ (} i = 1, \dots, k \text{)}. \end{aligned}$$

In Section 5 the following straightforward consequence of Lemma 4.1(i) will be useful.

Corollary 4.2. *Under the same assumptions as in Lemma 4.1, if e is an integer that is a multiple of the exponent of the group $\mathbf{M}_{\mathbf{C}}(\alpha, o)$, then*

$$(4.5) \quad d^{\mathbf{C}}(d^{\mathbf{C}}(\dots d^{\mathbf{C}}(d^{\mathbf{C}}(u_1, u_{e+1}, u_2), u_{e+1}, u_3) \dots), u_{e+1}, u_e) = u_1 +_o \dots +_o u_{e+1}$$

for all $u_1, \dots, u_{e+1} \in o^{\mathbf{C}}/\alpha$.

Proof. By Lemma 4.1(i) the left hand side of (4.5) equals $u_1 +_o u_2 +_o u_3 +_o \dots +_o u_e -_o (e-1)u_{e+1}$, which equals the right hand side of (4.5), because e is a multiple of the exponent of the group $\mathbf{M}_{\mathbf{C}}(\alpha, o)$. $\square$

Next we define the matrix ring $\mathbf{R}(\mathcal{V}, \mathcal{O})$ of the variety $\mathcal{V}$ (where the size of the matrices is $|\mathcal{O}|$). Let $\mathbf{F}$ be the free algebra in $\mathcal{V}$ with free generating set $\{x\} \cup \mathcal{O}$. In other words, $\mathbf{F}$ is the algebra of all $\mathcal{O}$ -terms of $\mathcal{V}_{\mathcal{O}}$ in one variable x (modulo the identities of $\mathcal{V}_{\mathcal{O}}$). For each $o \in \mathcal{O}$, let ε_o denote the unique endomorphism of $\mathbf{F}$ that maps x to o and fixes all elements of $\mathcal{O}$; that is, ε_o maps every $\mathcal{O}$ -term $r = r(x)$ to $r(o)$. Furthermore, let $\gamma_o := [\theta_o, \theta_o]$ where $\theta_o := \ker(\varepsilon_o)$. Now for any $o, o' \in \mathcal{O}$, let

$$\overline{H}_{o,o'} := \{r/\gamma_o : r = r(x) \in \mathbf{F}, r(o) = o'\}.$$

Then each $\overline{H}_{o,o'}$ has a natural abelian group structure with zero element o'/γ_o and addition and inversion defined by

$$\begin{aligned} r/\gamma_o + s/\gamma_o &:= d(r, o', s)/\gamma_o \quad \text{and} \\ -(r/\gamma_o) &:= d(o', r, o')/\gamma_o \quad \text{for all } r/\gamma_o, s/\gamma_o \in \overline{H}_{o,o'}. \end{aligned}$$

Moreover, composition of $\mathcal{O}$ -terms yields a multiplication that is defined by

$$(t/\gamma_{o'})(r/\gamma_o) := t(r(x))/\gamma_o \quad \text{for all } t/\gamma_{o'} \in \overline{H}_{o',o''} \text{ and } r/\gamma_o \in \overline{H}_{o,o'}.$$

It can be also shown that multiplication distributes over addition. Thus, the $\mathcal{O} \times \mathcal{O}$ matrices $(m_{o',o})$ with the properties that

- the entry $m_{o',o}$ in position (o', o) (i.e., row o' , column o) satisfies the condition $m_{o',o} \in \overline{H}_{o,o'}$ for every $o, o' \in \mathcal{O}$, and
- each column contains only finitely many nonzero entries

form a ring for the ordinary matrix operations. This is the matrix ring $\mathbf{R}(\mathcal{V}, \mathcal{O})$ associated to $\mathcal{V}$.

The proofs of all claims made throughout the definition above and the lemma below can be found at the beginning of Chapter 9 (before Theorem 9.4) in [10].

Lemma 4.3 (From [10]). *If α is an abelian congruence of an $\mathcal{O}$ -algebra $\mathbf{C}$ in $\mathcal{V}_{\mathcal{O}}$, then the direct sum*

$$\begin{aligned} \mathbf{M}_{\mathbf{C}}(\alpha, \mathcal{O}) &:= \bigoplus_{o \in \mathcal{O}} \mathbf{M}_{\mathbf{C}}(\alpha, o) \\ &= \{(a_o)_{o \in \mathcal{O}} \in \prod_{o \in \mathcal{O}} \mathbf{M}_{\mathbf{C}}(\alpha, o) : a_o = o^{\mathbf{C}} \text{ for all but finitely many } o\text{'s}\} \end{aligned}$$

of the abelian groups $\mathbf{M}_{\mathbf{C}}(\alpha, o)$ ($o \in \mathcal{O}$) is an $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -module for the ordinary matrix-vector multiplication if multiplication for the entries is defined as follows: for $a \in \mathbf{M}_{\mathbf{C}}(\alpha, o)$ and $m_{o', o} = r/\gamma_o \in \overline{H}_{o, o'}$ (with $r = r(x) \in \mathbf{F}$, $r(o) = o'$),

$$(4.6) \quad m_{o', o}a = (r/\gamma_o)a := r^{\mathbf{C}}(a).$$

Now we are ready to state our first theorem which, for every pair $(\mathbf{C}, \alpha)$ consisting of an $\mathcal{O}$ -algebra $\mathbf{C}$ in $\mathcal{V}_{\mathcal{O}}$ and an abelian congruence α of $\mathbf{C}$, relates the subalgebras of $\mathbf{C}$ to the submodules of the associated $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -module $\mathbf{M}_{\mathbf{C}}(\alpha, \mathcal{O})$. Recall our convention that the restriction of α to any subalgebra $\mathbf{U}$ (subset U) of $\mathbf{C}$ is denoted by $\alpha_{\mathbf{U}}$ (α_U , respectively).

Theorem 4.4. *Let α be an abelian congruence of an $\mathcal{O}$ -algebra $\mathbf{C}$ in $\mathcal{V}_{\mathcal{O}}$ such that the constants $o^{\mathbf{C}}$ ($o \in \mathcal{O}$) represent all α -classes of $\mathbf{C}$, and let $\mathbf{E} := \langle \mathcal{O}^{\mathbf{C}} \rangle$ be the least $\mathcal{O}$ -subalgebra of $\mathbf{C}$.*

- (1) *If $\mathbf{U}$ is an $\mathcal{O}$ -subalgebra of $\mathbf{C}$, then $\mathbf{M}_{\mathbf{U}}(\alpha_{\mathbf{U}}, \mathcal{O})$ is an $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -submodule of $\mathbf{M}_{\mathbf{C}}(\alpha, \mathcal{O})$ that contains $\mathbf{M}_{\mathbf{E}}(\alpha_{\mathbf{E}}, \mathcal{O})$ as a submodule.*
- (2) *The following conditions on any subset U of $\mathbf{C}$ are equivalent:*
 - (a) *U is the universe of a subalgebra of $\mathbf{C}$ that contains $\mathbf{E}$;*
 - (b) *U is the universe of an $\mathcal{O}$ -subalgebra of $\mathbf{C}$;*
 - (c) *U is closed under all functions $t^{\mathbf{C}}(x_1, \dots, x_k) \upharpoonright_{(o_1^{\mathbf{C}}/\alpha) \times \dots \times (o_k^{\mathbf{C}}/\alpha)}$ where t is an $\mathcal{O}$ -term and $o_1, \dots, o_k \in \mathcal{O}$;*
 - (d) *the set*

$$\bigoplus_{o \in \mathcal{O}} (o^{\mathbf{C}}/\alpha_U) := \{(u_o)_{o \in \mathcal{O}} \in \prod_{o \in \mathcal{O}} (o^{\mathbf{C}}/\alpha_U) : u_o = o^{\mathbf{C}} \text{ for all but finitely many } o\}$$

is the universe of an $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -submodule of $\mathbf{M}_{\mathbf{C}}(\alpha, \mathcal{O})$ that contains $\mathbf{M}_{\mathbf{E}}(\alpha_{\mathbf{E}}, \mathcal{O})$.

- (3) *If U is the universe of an $\mathcal{O}$ -subalgebra $\mathbf{U}$ of $\mathbf{C}$, then the $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -submodule of $\mathbf{M}_{\mathbf{C}}(\alpha, \mathcal{O})$ described in (d) is $\mathbf{M}_{\mathbf{U}}(\alpha_{\mathbf{U}}, \mathcal{O})$.*

Proof. For (1), assume that $\mathbf{U}$ is an $\mathcal{O}$ -subalgebra of $\mathbf{C}$; in particular, $o^{\mathbf{U}} = o^{\mathbf{C}}$ for all $o \in \mathcal{O}$. Then $\alpha_{\mathbf{U}}$ is an abelian congruence of $\mathbf{U}$, and the universe of $\mathbf{M}_{\mathbf{U}}(\alpha_{\mathbf{U}}, \mathcal{O})$ is a subset of the universe of $\mathbf{M}_{\mathbf{C}}(\alpha, \mathcal{O})$. The definition of the modules $\mathbf{M}_{\mathbf{U}}(\alpha_{\mathbf{U}}, \mathcal{O})$ and $\mathbf{M}_{\mathbf{C}}(\alpha, \mathcal{O})$ shows that they have the same zero element $(o^{\mathbf{U}})_{o \in \mathcal{O}} = (o^{\mathbf{C}})_{o \in \mathcal{O}}$, the group operations in the two modules are determined, in each coordinate, by the same $\mathcal{O}$ -terms $d(x, o, y)$, $d(o, x, o)$ (see Lemma 4.1), and for every matrix in $\mathbf{R}(\mathcal{V}, \mathcal{O})$, multiplication for the entries in the two modules are determined by the same $\mathcal{O}$ -terms (see Lemma 4.3). Since $\mathbf{U}$ is an $\mathcal{O}$ -subalgebra of $\mathbf{C}$, this implies that $\mathbf{M}_{\mathbf{U}}(\alpha_{\mathbf{U}}, \mathcal{O})$ is an $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -submodule of $\mathbf{M}_{\mathbf{C}}(\alpha, \mathcal{O})$.

$\mathbf{E}$ is an $\mathcal{O}$ -subalgebra of $\mathbf{U}$, therefore we can repeat the argument in the preceding paragraph for $\mathbf{U}$ and $\mathbf{E}$ (in place of $\mathbf{C}$ and $\mathbf{U}$) to conclude that $\mathbf{M}_{\mathbf{E}}(\alpha_{\mathbf{E}}, \mathcal{O})$ is an $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -submodule of $\mathbf{M}_{\mathbf{U}}(\alpha_{\mathbf{U}}, \mathcal{O})$.

To verify (2) and (3), we will prove the equivalence of the first three conditions in (2) by showing that (a) $\Leftrightarrow$ (b) and (b) $\Leftrightarrow$ (c). To establish that the fourth condition is also equivalent to them we will prove that (b) $\Rightarrow$ (d) and (d) $\Rightarrow$ (c). Statement (3) will be verified along with the proof of the implication (b) $\Rightarrow$ (d).

If U is the universe of a subalgebra $\mathbf{U}$ of $\mathbf{C}$ containing $\mathbf{E} = \langle \mathcal{O}^{\mathbf{C}} \rangle$, then we can define $o^{\mathbf{U}} := o^{\mathbf{C}}$ for every $o \in \mathcal{O}$ to make $\mathbf{U}$ an $\mathcal{O}$ -subalgebra of $\mathbf{C}$. This proves (a) $\Rightarrow$ (b). The converse is a tautology, so (a) $\Leftrightarrow$ (b) is proved.

U is the universe of an $\mathcal{O}$ -subalgebra of $\mathbf{C}$ if and only if U is closed under all functions $t^{\mathbf{C}}$ induced by $\mathcal{O}$ -terms t . Since our assumption is that the constants $o^{\mathbf{C}}$ ($o \in \mathcal{O}$) represent all α -classes of $\mathbf{C}$, it follows that for every $\mathcal{O}$ -term t , U is closed under the term function $t^{\mathbf{C}}$ if and only if it is closed under all its restrictions described in (c). This proves (b) $\Leftrightarrow$ (c).

Now assume that U is the universe of an $\mathcal{O}$ -subalgebra $\mathbf{U}$ of $\mathbf{C}$; in particular, $o^{\mathbf{U}} = o^{\mathbf{C}}$ for all $o \in \mathcal{O}$. Then $\alpha_U = \alpha_{\mathbf{U}}$ is an abelian congruence of $\mathbf{U}$. The definition of the $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -module $\mathbf{M}_{\mathbf{U}}(\alpha_{\mathbf{U}}, \mathcal{O})$ shows that its universe is exactly $\bigoplus_{o \in \mathcal{O}} (o^{\mathbf{C}} / \alpha_U)$. Therefore statement (1) proves that $\bigoplus_{o \in \mathcal{O}} (o^{\mathbf{C}} / \alpha_U)$ is indeed the underlying set of an $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -submodule of $\mathbf{M}_{\mathbf{C}}(\alpha, \mathcal{O})$ that contains $\mathbf{M}_{\mathbf{E}}(\alpha_{\mathbf{E}}, \mathcal{O})$ as a submodule, namely $\mathbf{M}_{\mathbf{U}}(\alpha_{\mathbf{U}}, \mathcal{O})$. This finishes the proof of both (b) $\Rightarrow$ (d) and (3).

Finally, we want to argue that (d) $\Rightarrow$ (c). We will start by proving a claim which is independent of U .

Claim 4.5. *Let $\mathbf{C}$ and α be as in the theorem. For arbitrary $\mathcal{O}$ -term $t = t(x_1, \dots, x_k)$ and constant symbols $o_1, \dots, o_k \in \mathcal{O}$,*

- (1) $t^{\mathbf{C}}(o_1^{\mathbf{C}}, \dots, o_k^{\mathbf{C}}) \in o^{\mathbf{C}} / \alpha$ for some $o \in \mathcal{O}$, and
- (2) *for every such $o \in \mathcal{O}$ there exist $m_i = r_i / \gamma_{o_i} \in \overline{H}_{o_i, o}$ (with $r_i = r_i(x) \in \mathbf{F}$, $r_i(o_i) = o$) for $i = 1, \dots, k$ such that*

$$(4.7) \quad t^{\mathbf{C}}(u_1, \dots, u_k) = m_1 u_1 +_o \dots +_o m_k u_k +_o t^{\mathbf{C}}(o_1^{\mathbf{C}}, \dots, o_k^{\mathbf{C}})$$

for all $(u_1, \dots, u_k) \in (o_1^{\mathbf{C}} / \alpha) \times \dots \times (o_k^{\mathbf{C}} / \alpha)$.

Proof of Claim 4.5. To simplify notation, let $\overline{x} := (x_1, \dots, x_k)$ and $\overline{o} := (o_1, \dots, o_k)$.

(1) holds, because our assumption that the constants $o^{\mathbf{C}}$ ($o \in \mathcal{O}$) represent all α -classes of $\mathbf{C}$ implies that the element $t^{\mathbf{C}}(\overline{o}^{\mathbf{C}})$ of $\mathbf{C}$ is in $o^{\mathbf{C}} / \alpha$ for some $o \in \mathcal{O}$.

To prove (2) let us fix such an $o \in \mathcal{O}$ for the rest of the argument. Since $t = t(\overline{x})$ is an $\mathcal{O}$ -term, so is

$$(4.8) \quad r(\overline{x}) := d(t(\overline{x}), t(\overline{o}), o).$$

As in Lemma 4.1(ii), consider the unary $\mathcal{O}$ -terms $r_i(x) := r(o_1, \dots, o_{i-1}, x, o_{i+1}, \dots, o_k)$ for each i ($1 \leq i \leq k$). Since d satisfies the identity $d(x, x, y) = y$ (see (4.1)), we get that $r(\overline{o}) = o$, and hence $r_i(o_i) = o$ for every i ($1 \leq i \leq k$). This shows that r satisfies the assumptions of Lemma 4.1(ii), and $m_i := r_i / \gamma_{o_i}$ is a member of $\overline{H}_{o_i, o}$ for

every i ($1 \leq i \leq k$). Thus, first using equality (4.4) from Lemma 4.1(ii), and then equality (4.6) from Lemma 4.3 we obtain that

$$(4.9) \quad r^{\mathbf{C}}(u_1, \dots, u_k) = r_1^{\mathbf{C}}(u_1) +_o \dots +_o r_k^{\mathbf{C}}(u_k) = m_1 u_1 +_o \dots +_o m_k u_k$$

$$\text{for all } (u_1, \dots, u_k) \in (o_1^{\mathbf{C}}/\alpha) \times \dots \times (o_k^{\mathbf{C}}/\alpha).$$

On the other hand, since $r^{\mathbf{C}}(u_1, \dots, u_k) \in o^{\mathbf{C}}/\alpha$ for all $(u_1, \dots, u_k) \in (o_1^{\mathbf{C}}/\alpha) \times \dots \times (o_k^{\mathbf{C}}/\alpha)$ (cf. Lemma 4.1(ii)), we can apply first (4.8), and then the equality in Lemma 4.1(i) to get that

$$r^{\mathbf{C}}(u_1, \dots, u_k) = d^{\mathbf{C}}(t^{\mathbf{C}}(u_1, \dots, u_k), t^{\mathbf{C}}(\bar{o}^{\mathbf{C}}), o^{\mathbf{C}}) = t^{\mathbf{C}}(u_1, \dots, u_k) -_o t^{\mathbf{C}}(\bar{o}^{\mathbf{C}})$$

$$\text{for all } (u_1, \dots, u_k) \in (o_1^{\mathbf{C}}/\alpha) \times \dots \times (o_k^{\mathbf{C}}/\alpha).$$

The last displayed equality shows that by adding $t^{\mathbf{C}}(\bar{o}^{\mathbf{C}})$ to both sides of (4.9) in the group $\mathbf{M}_{\mathbf{C}}(\alpha, o)$ we get the equality (4.7) we wanted to prove. $\diamond$

To prove the implication (d) $\Rightarrow$ (c), assume that (d) holds for U , that is, the subset $S_U := \bigoplus_{o \in \mathcal{O}} (o^{\mathbf{C}}/\alpha_U)$ of $\mathbf{M}_{\mathbf{C}}(\alpha, \mathcal{O})$ is the universe of an $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -submodule of $\mathbf{M}_{\mathbf{C}}(\alpha, \mathcal{O})$, and contains all elements of $\mathbf{M}_{\mathbf{E}}(\alpha_{\mathbf{E}}, \mathcal{O})$. In particular, the zero element $(o^{\mathbf{C}})_{o \in \mathcal{O}}$ of $\mathbf{M}_{\mathbf{C}}(\alpha, \mathcal{O})$ belongs to S_U , which implies that $\mathcal{O}^{\mathbf{C}} \subseteq U$.

Let $t = t(x_1, \dots, x_k)$ be an arbitrary $\mathcal{O}$ -term, and let $o_1, \dots, o_k \in \mathcal{O}$. We want to show that U is closed under the function $t^{\mathbf{C}}(x_1, \dots, x_k) \upharpoonright_{(o_1^{\mathbf{C}}/\alpha) \times \dots \times (o_k^{\mathbf{C}}/\alpha)}$. By Claim 4.5 there exist $o \in \mathcal{O}$ and $m_i = r_i/\gamma_{o_i} \in \overline{H}_{o_i, o}$ (with $r_i = r_i(x) \in \mathbf{F}$, $r_i(o_i) = o$) for $i = 1, \dots, k$ such that $t^{\mathbf{C}}(o_1^{\mathbf{C}}, \dots, o_k^{\mathbf{C}}) \in o^{\mathbf{C}}/\alpha$ and (4.7) holds. Let us fix an arbitrary tuple $(u_1, \dots, u_k) \in (o_1^{\mathbf{C}}/\alpha) \times \dots \times (o_k^{\mathbf{C}}/\alpha)$ that belongs to U ; that is, $(u_1, \dots, u_k) \in (o_1^{\mathbf{C}}/\alpha_U) \times \dots \times (o_k^{\mathbf{C}}/\alpha_U)$. Furthermore, let $a := t^{\mathbf{C}}(o_1^{\mathbf{C}}, \dots, o_k^{\mathbf{C}})$. We will be done if we show that the element

$$b := t^{\mathbf{C}}(u_1, \dots, u_k) = m_1 u_1 +_o \dots +_o m_k u_k +_o a$$

belongs to U .

Note that $b = t^{\mathbf{C}}(u_1, \dots, u_k) \equiv_{\alpha} t^{\mathbf{C}}(o_1^{\mathbf{C}}, \dots, o_k^{\mathbf{C}}) = a$, therefore $a \in o^{\mathbf{C}}/\alpha$ implies that $b \in o^{\mathbf{C}}/\alpha$. Moreover, since $a = t^{\mathbf{C}}(o_1^{\mathbf{C}}, \dots, o_k^{\mathbf{C}})$ is an element of the least $\mathcal{O}$ -subalgebra $\mathbf{E}$ of $\mathbf{C}$ (in which $o^{\mathbf{E}} = o^{\mathbf{C}}$), we get from $a \in o^{\mathbf{C}}/\alpha$ that $a \in o^{\mathbf{E}}/\alpha_{\mathbf{E}}$.

Now, for each i ($1 \leq i \leq k$), let $\tilde{m}_i$ denote the $\mathcal{O} \times \mathcal{O}$ matrix which has m_i in position (o, o_i) and zeros (i.e., $o''/\gamma_{o'}$) in all other positions (o'', o') , and let $\tilde{u}_i$ denote the $\mathcal{O}$ -tuple which has u_i in its o_i -th position and zeros (i.e., $o'^{\mathbf{C}}$) in all other positions $o' \in \mathcal{O}$. Furthermore, let $\tilde{a}$ and $\tilde{b}$ denote the $\mathcal{O}$ -tuples which have a and b , respectively, in their o -th positions, and zeros in all other positions. Since $m_i \in \overline{H}_{o_i, o}$ for every i ($1 \leq i \leq k$), the definition of $\mathbf{R}(\mathcal{V}, \mathcal{O})$ shows that $\tilde{m}_1, \dots, \tilde{m}_k \in \mathbf{R}(\mathcal{V}, \mathcal{O})$. Since $u_i \in o_i^{\mathbf{C}}/\alpha_U$ for every i ($1 \leq i \leq k$) and $\mathcal{O}^{\mathbf{C}} \subseteq U$, we get that $\tilde{u}_1, \dots, \tilde{u}_k \in S_U \subseteq \mathbf{M}_{\mathbf{C}}(\alpha, \mathcal{O})$. Similarly, $a \in o^{\mathbf{E}}/\alpha_{\mathbf{E}}$ implies that $\tilde{a} \in \mathbf{M}_{\mathbf{E}}(\alpha_{\mathbf{E}}, \mathcal{O})$. Therefore, our assumption that S_U contains the elements of $\mathbf{M}_{\mathbf{E}}(\alpha_{\mathbf{E}}, \mathcal{O})$ yields that $\tilde{a} \in S_U \subseteq \mathbf{M}_{\mathbf{C}}(\alpha, \mathcal{O})$. Finally,

we have $\tilde{b} \in \mathbf{M}_{\mathbf{C}}(\alpha, \mathcal{O})$, because $b \in o^{\mathbf{C}}/\alpha$. Since $b = m_1 u_1 +_o \cdots +_o m_k u_k +_o a$, the construction of the tuples $\tilde{u}_i$, $\tilde{a}$, $\tilde{b}$ and the matrices $\tilde{m}_i$ makes sure that the equality $\tilde{b} = \tilde{m}_1 \tilde{u}_1 + \cdots + \tilde{m}_k \tilde{u}_k + \tilde{a}$ holds in the $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -module $\mathbf{M}_{\mathbf{C}}(\alpha, \mathcal{O})$. Since $\tilde{u}_1, \dots, \tilde{u}_k, \tilde{a} \in S_U$ and, by our assumption, S_U is closed under the $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -module operations of $\mathbf{M}_{\mathbf{C}}(\alpha, \mathcal{O})$, we get that $\tilde{b} \in S_U$. Hence, $b \in o^{\mathbf{C}}/\alpha_U \subseteq U$. This completes the proof of (d) $\Rightarrow$ (c), and also the proof of Theorem 4.4 $\square$

Our second theorem is the analog of Theorem 9.9 in [10] mentioned at the beginning of this section. Given a pair $(\mathbf{C}, \alpha)$ consisting of an $\mathcal{O}$ -algebra in $\mathcal{V}_{\mathcal{O}}$ and an abelian congruence α of $\mathbf{C}$, the theorem establishes an isomorphism between the lattice of $\mathcal{O}$ -subalgebras of $\mathbf{C}$ and an interval in the submodule lattice of the associated $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -module $\mathbf{M}_{\mathbf{C}}(\alpha, \mathcal{O})$, provided all α -classes are represented in the module.

Theorem 4.6. *Let $\mathbf{C}$ be an $\mathcal{O}$ -algebra in $\mathcal{V}_{\mathcal{O}}$, and let $\mathbf{E} := \langle \mathcal{O}^{\mathbf{C}} \rangle$ be its least $\mathcal{O}$ -subalgebra. If α is an abelian congruence of $\mathbf{C}$ such that the constants $o^{\mathbf{C}}$ ($o \in \mathcal{O}$) represent all α -classes of $\mathbf{C}$, then the mapping*

$$(4.10) \quad \mathbf{U} \mapsto \mathbf{M}_{\mathbf{U}}(\alpha_{\mathbf{U}}, \mathcal{O})$$

is an isomorphism between the lattice of all $\mathcal{O}$ -subalgebras of $\mathbf{C}$ and the lattice of all $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -submodules of $\mathbf{M}_{\mathbf{C}}(\alpha, \mathcal{O})$ that contain $\mathbf{M}_{\mathbf{E}}(\alpha_{\mathbf{E}}, \mathcal{O})$.

Proof. It follows from Theorem 4.4(1) that for every $\mathcal{O}$ -subalgebra $\mathbf{U}$ of $\mathbf{C}$, $\mathbf{M}_{\mathbf{U}}(\alpha_{\mathbf{U}}, \mathcal{O})$ is an $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -submodule of $\mathbf{M}_{\mathbf{C}}(\alpha, \mathcal{O})$ that contains $\mathbf{M}_{\mathbf{E}}(\alpha_{\mathbf{E}}, \mathcal{O})$. Let $\mathbf{U}$ and $\mathbf{V}$ be arbitrary $\mathcal{O}$ -subalgebras of $\mathbf{C}$. If $\mathbf{U} \leq \mathbf{V}$, then for every $o \in \mathcal{O}$, the congruence class $o^{\mathbf{U}}/\alpha_{\mathbf{U}} = o^{\mathbf{C}}/\alpha_{\mathbf{U}}$ in $\mathbf{U}$ is contained in the corresponding congruence class $o^{\mathbf{V}}/\alpha_{\mathbf{V}} = o^{\mathbf{C}}/\alpha_{\mathbf{V}}$ of $\mathbf{V}$. This implies that $\mathbf{M}_{\mathbf{U}}(\alpha_{\mathbf{U}}, \mathcal{O}) \leq \mathbf{M}_{\mathbf{V}}(\alpha_{\mathbf{V}}, \mathcal{O})$, showing that the map (4.10) is order-preserving. Now suppose that $\mathbf{U} \not\leq \mathbf{V}$. The assumption that the interpretations of the constant symbols $o \in \mathcal{O}$ represent all α -classes of $\mathbf{C}$, implies the existence of $o^{\mathbf{C}} = o^{\mathbf{U}} = o^{\mathbf{V}}$ such that $o^{\mathbf{U}}/\alpha_{\mathbf{U}} \not\subseteq o^{\mathbf{V}}/\alpha_{\mathbf{V}}$. Thus, $\mathbf{M}_{\mathbf{U}}(\alpha_{\mathbf{U}}, \mathcal{O}) \not\leq \mathbf{M}_{\mathbf{V}}(\alpha_{\mathbf{V}}, \mathcal{O})$. This shows that the map (4.10) preserves and reflects $\leq$; hence, in particular, it is one-to-one.

The surjectivity of the map (4.10) is proved by the next claim.

Claim 4.7. *Let $\mathbf{C}$, $\mathbf{E}$, and α be as in Theorem 4.6. If $\mathbf{N}$ is an $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -submodule of $\mathbf{M}_{\mathbf{C}}(\alpha, \mathcal{O})$ that contains $\mathbf{M}_{\mathbf{E}}(\alpha_{\mathbf{E}}, \mathcal{O})$, then*

- (1) *as an abelian group, $\mathbf{N} = \bigoplus_{o \in \mathcal{O}} \mathbf{N}_o$ where for every $o \in \mathcal{O}$, $\mathbf{N}_o$ is the subgroup of $\mathbf{M}_{\mathbf{C}}(\alpha, o)$ that consists of the o -components of all tuples in $\mathbf{N}$;*
- (2) *$\mathbf{N}_o = \mathbf{N}_{o'}$ whenever $o^{\mathbf{C}}/\alpha = o'^{\mathbf{C}}/\alpha$ ($o, o' \in \mathcal{O}$);*
- (3) *the union $U := \bigcup_{o \in \mathcal{O}} \mathbf{N}_o$ of the universes of the groups $\mathbf{N}_o$ is the universe of an $\mathcal{O}$ -subalgebra $\mathbf{U}$ of $\mathbf{C}$, and $\mathbf{N} = \mathbf{M}_{\mathbf{U}}(\alpha_{\mathbf{U}}, \mathcal{O})$.*

Proof of Claim 4.7. For (1) notice first that since the group operations in $\mathbf{M}_{\mathbf{C}}(\alpha, \mathcal{O})$ are performed coordinatewise, it follows immediately from the definition of the $\mathbf{N}_o$'s

that $\mathbf{N}_o$ is a subgroup of $\mathbf{M}_C(\alpha, o)$ (in particular, $o^C \in \mathbf{N}_o$) for each $o \in \mathcal{O}$, and $\mathbf{N} \subseteq \bigoplus_{o \in \mathcal{O}} \mathbf{N}_o$.

To prove that equality holds, consider an arbitrary element $(a_o)_{o \in \mathcal{O}} \in \bigoplus_{o \in \mathcal{O}} \mathbf{N}_o (\subseteq \mathbf{M}_C(\alpha, \mathcal{O}))$. Then (i) $a_o \in N_o$ for every $o \in \mathcal{O}$, and (ii) $a_o = o^C$ for all but finitely many o 's. Condition (i) implies that for every $o \in \mathcal{O}$ there exists a tuple $\bar{a}_o \in \mathbf{N}$ such that the o -th coordinate of $\bar{a}_o$ is a_o . Let e_o be the matrix in $\mathbf{R}(\mathcal{V}, \mathcal{O})$ which has x/γ_o in position (o, o) , and zeros (that is, $o''/\gamma_{o'}$) in all other positions (o'', o') . Since $\mathbf{N}$ is an $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -submodule of $\mathbf{M}_C(\alpha, \mathcal{O})$, each tuple $e_o \bar{a}_o$ is an element of $\mathbf{N}$. It follows from the construction of e_o that the tuple $e_o \bar{a}_o$ has a_o in the o -th coordinate and zeros elsewhere. By condition (ii), only finitely many of the tuples $e_o \bar{a}_o$ ($o \in \mathcal{O}$) is not the zero element of $\mathbf{N}$, therefore their sum in $\mathbf{M}_C(\alpha, \mathcal{O})$, and hence also in $\mathbf{N}$, is $(a_o)_{o \in \mathcal{O}}$. Thus $(a_o)_{o \in \mathcal{O}} \in \mathbf{N}$, which finishes the proof that $\mathbf{N} = \bigoplus_{o \in \mathcal{O}} \mathbf{N}_o$.

To prove (2) assume that $o, o' \in \mathcal{O}$ are such that $o^C/\alpha = o'^C/\alpha$. Since $o^E = o^C$ and $o'^E = o'^C$, we also have that $o^E/\alpha_E = o'^E/\alpha_E$. Therefore our assumption that $\mathbf{N}$ is an $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -submodule of $\mathbf{M}_C(\alpha, \mathcal{O})$ that contains $\mathbf{M}_E(\alpha_E, \mathcal{O})$ implies that

$$o^E/\alpha_E = o'^E/\alpha_E \subseteq N_o \cap N_{o'} \subseteq N_o \cup N_{o'} \subseteq o^C/\alpha = o'^C/\alpha.$$

In particular, we have that $o^E, o'^E \in N_o \cap N_{o'}$, or equivalently, $o^C, o'^C \in N_o \cap N_{o'}$.

For any element $a \in N_o$ let $\tilde{a}$ denote the $\mathcal{O}$ -tuple which has a in its o -th position, and zeros in all other positions (i.e., o''^C in the o'' -th position for $o'' \neq o$). It follows from statement (1) that $\tilde{a} \in \mathbf{N}$ for all $a \in N_o$. Now let r denote the $\mathcal{O} \times \mathcal{O}$ matrix which has $d(x, o, o')/\gamma_o$ in position (o', o) and zeros (i.e., $o'''/\gamma_{o''}$) in all other positions (o''', o'') . Since $d(o, o, o') = o'$, we have that $d(x, o, o')/\gamma_o \in \overline{H}_{o, o'}$, and hence $r \in \mathbf{R}(\mathcal{V}, \mathcal{O})$. As $\mathbf{N}$ is an $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -submodule of $\mathbf{M}_C(\alpha, \mathcal{O})$, we get that $r\tilde{a} \in \mathbf{N}$ for all $a \in N_o$. Thus the o' -th component of $r\tilde{a}$ is in $N_{o'}$; that is,

$$(4.11) \quad d^C(a, o^C, o'^C) = (d(x, o, o')/\gamma_o)a \in N_{o'} \quad \text{for all } a \in N_o.$$

By Lemma 4.1(i), $d^C(u, o^C, o'^C) = u +_o o'^C$ holds in $\mathbf{M}_C(\alpha, o)$ for all $u \in o^C/\alpha$. Therefore (4.11) can be restated as follows: the image of N_o under the translation $u \mapsto u +_o o'^C$ is contained in $N_{o'}$. Because of $o'^C \in N_o$ this translation is a bijection of the abelian group $\mathbf{N}_o$ onto itself, so we get that $N_o \subseteq N_{o'}$. A similar argument, with the roles of o and o' switched, shows that $N_{o'} \subseteq N_o$ also holds, and hence finishes the proof of statement (2).

Finally, to establish (3), recall that $U := \bigcup_{o \in \mathcal{O}} N_o$. Since $(o^C \in) N_o \subseteq o^C/\alpha (\subseteq C)$ for every $o \in \mathcal{O}$, we get that $U \subseteq C$ and that the sets N_o and $N_{o'}$ are disjoint whenever $o^C \not\equiv_\alpha o'^C$. By statement (2), $N_o = N_{o'}$ whenever $o^C \equiv_\alpha o'^C$. Therefore $\{N_o : o \in \mathcal{O}\}$ is the partition of U corresponding to the equivalence relation α_U . It follows now from statement (1) that the universe of $\mathbf{N}$ is the set $\bigoplus_{o \in \mathcal{O}} N_o = \bigoplus_{o \in \mathcal{O}} (o^C/\alpha_U)$. Hence, the equivalence of conditions (d) and (b) in Theorem 4.4(2) implies that U

is the universe of an $\mathcal{O}$ -subalgebra $\mathbf{U}$ of $\mathbf{C}$. Moreover, by Theorem 4.4(3), we have that $\mathbf{N} = \mathbf{M}_{\mathbf{U}}(\alpha_{\mathbf{U}}, \mathcal{O})$. This finishes the proof of Claim 4.7. $\diamond$

The proof of Theorem 4.6 is complete. $\square$

We conclude this section by an auxiliary result concerning the modules associated to α -saturated subalgebras of direct products where α is an abelian product congruence.

Lemma 4.8. *Let $\mathbf{C}$ be an $\mathcal{O}$ -subalgebra of an $\mathcal{O}$ -product $\prod_{i=1}^n \mathbf{B}_i$ of $\mathcal{O}$ -algebras $\mathbf{B}_1, \dots, \mathbf{B}_n$ in $\mathcal{V}_{\mathcal{O}}$, and let $\alpha = \prod_{i=1}^n \alpha_i$ be the product congruence of $\prod_{i=1}^n \mathbf{B}_i$ where for each i , α_i is an abelian congruence of $\mathbf{B}_i$. If $\mathbf{C}$ is α -saturated in $\prod_{i=1}^n \mathbf{B}_i$, then*

- (1) $\alpha_{\mathbf{C}}$ is an abelian congruence of $\mathbf{C}$ and $o^{\mathbf{C}}/\alpha_{\mathbf{C}} = \prod_{i=1}^n (o^{\mathbf{B}_i}/\alpha_i)$ holds for all $o \in \mathcal{O}$; moreover,
- (2) the $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -modules $\mathbf{M}_{\mathbf{C}}(\alpha_{\mathbf{C}}, \mathcal{O})$ and $\prod_{i=1}^n \mathbf{M}_{\mathbf{B}_i}(\alpha_i, \mathcal{O})$ are naturally isomorphic via the map defined by

$$(4.12) \quad \mathbf{M}_{\mathbf{C}}(\alpha_{\mathbf{C}}, \mathcal{O}) \rightarrow \prod_{i=1}^n \mathbf{M}_{\mathbf{B}_i}(\alpha_i, \mathcal{O})$$

$$(a_o)_{o \in \mathcal{O}} = ((a_{o1}, \dots, a_{on}))_{o \in \mathcal{O}} \mapsto ((a_{o1})_{o \in \mathcal{O}}, \dots, (a_{on})_{o \in \mathcal{O}})$$

for all $a_o = (a_{o1}, \dots, a_{on}) \in o^{\mathbf{C}}/\alpha_{\mathbf{C}} = \prod_{i=1}^n (o^{\mathbf{B}_i}/\alpha_i)$ ($o \in \mathcal{O}$).

Proof. First we prove (1). Since each α_i is an abelian congruence of $\mathbf{B}_i$, their product, α , is an abelian congruence of $\prod_{i=1}^n \mathbf{B}_i$, and hence $\alpha_{\mathbf{C}}$ is an abelian congruence of $\mathbf{C}$. Therefore the module $\mathbf{M}_{\mathbf{C}}(\alpha_{\mathbf{C}}, \mathcal{O})$ exists. By our assumption, $\prod_{i=1}^n \mathbf{B}_i$ is an $\mathcal{O}$ -product and $\mathbf{C}$ is an $\mathcal{O}$ -subalgebra of $\prod_{i=1}^n \mathbf{B}_i$, so $o^{\mathbf{C}} = (o^{\mathbf{B}_1}, \dots, o^{\mathbf{B}_n})$ for every $o \in \mathcal{O}$. This implies that $o^{\mathbf{C}}/\alpha_{\mathbf{C}} \subseteq \prod_{i=1}^n (o^{\mathbf{B}_i}/\alpha_i)$ for every $o \in \mathcal{O}$. In fact, $=$ holds here, because $\mathbf{C}$ is α -saturated in $\prod_{i=1}^n \mathbf{B}_i$.

Now we prove (2). It follows from the equalities in (1) that the elements of $\mathbf{M}_{\mathbf{C}}(\alpha_{\mathbf{C}}, \mathcal{O})$ are exactly the tuples $(a_o)_{o \in \mathcal{O}}$ such that (i) $a_o \in o^{\mathbf{C}}/\alpha_{\mathbf{C}} = \prod_{i=1}^n (o^{\mathbf{B}_i}/\alpha_i)$, that is, $a_o = (a_{o1}, \dots, a_{on})$ with $a_{oi} \in o^{\mathbf{B}_i}/\alpha_i$ for every o and i , and (ii) $a_o = o^{\mathbf{C}} = (o^{\mathbf{B}_1}, \dots, o^{\mathbf{B}_n})$ for all but finitely many o 's. The image of each such tuple $(a_o)_{o \in \mathcal{O}}$ under the map (4.12) clearly belongs to the module $\prod_{i=1}^n \mathbf{M}_{\mathbf{B}_i}(\alpha_i, \mathcal{O})$. The map (4.12) just regroups coordinates, so it is one-to-one. To see that it is also onto, let $((b_{o1})_{o \in \mathcal{O}}, \dots, (b_{on})_{o \in \mathcal{O}}) \in \prod_{i=1}^n \mathbf{M}_{\mathbf{B}_i}(\alpha_i, \mathcal{O})$. Then, for every i , we have that (i)' $b_{oi} \in o^{\mathbf{B}_i}/\alpha_i$ for all $o \in \mathcal{O}$, and (ii)' $b_{oi} = o^{\mathbf{B}_i}$ for all but finitely many o 's. Therefore the tuples $b_o := (b_{o1}, \dots, b_{on})$ satisfy the conditions that (i) $b_o \in \prod_{i=1}^n (o^{\mathbf{B}_i}/\alpha_i) = o^{\mathbf{C}}/\alpha_{\mathbf{C}}$ for all $o \in \mathcal{O}$, and (ii) $b_o = (o^{\mathbf{B}_1}, \dots, o^{\mathbf{B}_n}) = o^{\mathbf{C}}$ for all but finitely many o 's. Hence $(b_o)_{o \in \mathcal{O}} \in \mathbf{M}_{\mathbf{C}}(\alpha_{\mathbf{C}}, \mathcal{O})$, and $((b_{o1})_{o \in \mathcal{O}}, \dots, (b_{on})_{o \in \mathcal{O}})$ is its image under the map (4.12). This proves that (4.12) is a bijection.

To see that (4.12) is a group isomorphism, recall from Lemma 4.3 that in each one of the modules $\mathbf{M}_{\mathbf{C}}(\alpha_{\mathbf{C}}, \mathcal{O})$ and $\mathbf{M}_{\mathbf{B}_i}(\alpha_i, \mathcal{O})$ ($1 \leq i \leq n$), $+$ is defined to be the

coordinatewise operation that is $+_o$ in coordinate o for each $o \in \mathcal{O}$. Therefore, in the module $\prod_{i=1}^n \mathbf{M}_{\mathbf{B}_i}(\alpha_i, \mathcal{O})$, $+$ is the coordinatewise operation that is the operation $+_o$ on $o^{\mathbf{B}_i}/\alpha_i (\subseteq \mathbf{B}_i)$ in coordinate (o, i) . In the module $\mathbf{M}_{\mathbf{C}}(\alpha_{\mathbf{C}}, \mathcal{O})$, the operation $+_o$ on $o^{\mathbf{C}}/\alpha_{\mathbf{C}} = \prod_{i=1}^n (o^{\mathbf{B}_i}/\alpha_i)$ also acts coordinatewise, because $+_o$ is an $\mathcal{O}$ -term and $\mathbf{C}$ is an $\mathcal{O}$ -subalgebra of $\prod_{i=1}^n \mathbf{B}_i$. Therefore, in $\mathbf{M}_{\mathbf{C}}(\alpha_{\mathbf{C}}, \mathcal{O})$, too, $+$ is the coordinatewise operation that is the operation $+_o$ on $o^{\mathbf{B}_i}/\alpha_i (\subseteq \mathbf{B}_i)$ in coordinate (o, i) . This shows that (4.12) is a group isomorphism.

Finally, we will argue that (4.12) is an $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -module isomorphism. Recall again from Lemma 4.3 and the discussion preceding it that $\mathbf{R}(\mathcal{V}, \mathcal{O})$ is a ring of matrices, and multiplication by elements of $\mathbf{R}(\mathcal{V}, \mathcal{O})$ in each one of the modules $\mathbf{M}_{\mathbf{C}}(\alpha_{\mathbf{C}}, \mathcal{O})$ and $\mathbf{M}_{\mathbf{B}_i}(\alpha_i, \mathcal{O})$ ($1 \leq i \leq n$) is defined by matrix multiplication. Furthermore, in the module $\prod_{i=1}^n \mathbf{M}_{\mathbf{B}_i}(\alpha_i, \mathcal{O})$, multiplication by ring elements is performed coordinatewise. Therefore we will be done if we show that

- (*) multiplication by ring elements acts coordinatewise on the components $a_o = (a_{o1}, \dots, a_{on})$ of the elements $(a_o)_{o \in \mathcal{O}}$ of $\mathbf{M}_{\mathbf{C}}(\alpha_{\mathbf{C}}, \mathcal{O})$.

Since (4.12) is a group isomorphism, it suffices to prove (*) for the multiplication of entries. Following the definition of multiplication of entries in Lemma 4.3, let $m_{o', o} = r/\gamma_o \in \overline{H}_{o, o'}$ (where $r = r(x) \in \mathbf{F}$ and $r(o) = o'$), and let $a_o = (a_{o1}, \dots, a_{on}) \in o^{\mathbf{C}}/\alpha_{\mathbf{C}} = \prod_{i=1}^n (o^{\mathbf{B}_i}/\alpha_i)$. By definition, $m_{o', o} a_o = r^{\mathbf{C}}(a_o)$. Since r is an $\mathcal{O}$ -term and $\mathbf{C}$ is an $\mathcal{O}$ -subalgebra of $\prod_{i=1}^n \mathbf{B}_i$, we get that $r^{\mathbf{C}}(a_o) = (r^{\mathbf{B}_i}(a_{oi}))_{i=1}^n = (m_{o', o} a_{oi})_{i=1}^n$. This proves that multiplication by $m_{o', o}$ acts coordinatewise on a_o , and hence finishes the proof of the lemma. $\square$

The isomorphism (4.12) shows that under the assumptions of Lemma 4.8 the only difference between the modules $\mathbf{M}_{\mathbf{C}}(\alpha_{\mathbf{C}}, \mathcal{O})$ and $\prod_{i=1}^n \mathbf{M}_{\mathbf{B}_i}(\alpha_i, \mathcal{O})$ is how we group their coordinates. Therefore we may identify $\mathbf{M}_{\mathbf{C}}(\alpha_{\mathbf{C}}, \mathcal{O})$ and $\prod_{i=1}^n \mathbf{M}_{\mathbf{B}_i}(\alpha_i, \mathcal{O})$ via the isomorphism (4.12), and view the submodules of $\mathbf{M}_{\mathbf{C}}(\alpha_{\mathbf{C}}, \mathcal{O})$ as submodules of the direct product $\prod_{i=1}^n \mathbf{M}_{\mathbf{B}_i}(\alpha_i, \mathcal{O})$.

5. PROOF OF THEOREM 1.1

Throughout this section $\mathbf{A}$ will be a finite algebra with a k -parallelogram term, and $\mathcal{V}$ the variety generated by $\mathbf{A}$. Our goal is to combine the results of Sections 3 and 4 to prove that if $\mathbf{A}$ satisfies the split centralizer condition, then $\mathbf{A}$ is dualizable. In view of Theorem 2.4, we will be done if we can show that there exists a constant c , depending only on $\mathbf{A}$, such that

$$(5.1) \quad \mathcal{R}_c(\mathbf{A}) \models_d \mathcal{R}(\mathbf{A}).$$

Recall that $\mathcal{R}(\mathbf{A})$ stands for the set of all (finitary) compatible relations of $\mathbf{A}$, and for every positive integer n , $\mathcal{R}_n(\mathbf{A})$ denotes the set of all compatible relations of $\mathbf{A}$ of arity $\leq n$. The relation $\models_d$ is described in Theorem 2.3.

In addition to the natural numbers $\mathbf{a}$, $\mathbf{s}$, $\mathbf{i}$, and $\mathbf{p}$ introduced in Section 3, we will need a few other parameters related to $\mathbf{A}$, which we introduce now. The list includes the constant $\mathbf{c}$ that we will use in the proof of (5.1). Before defining the new parameters, let us fix a set $\mathcal{O}$ of constant symbols not occurring in the language of $\mathcal{V}$ such that $|\mathcal{O}| = \mathbf{i}$. Furthermore, let $\text{Mod}(\mathbf{A}, \mathbf{p})$ denote the set of all $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -modules $\mathbf{M}_{\mathbf{T}}(\alpha, \mathcal{O})$ where $\mathbf{T}$ is a subalgebra of $\mathbf{A}^p$ for some $1 \leq p \leq \mathbf{p}$ that is isomorphic to a section of $\mathbf{A}$, α is a nontrivial abelian congruence of $\mathbf{T}$, and $\mathbf{T}$ is made into an $\mathcal{O}$ -algebra by interpreting the new constant symbols $o \in \mathcal{O}$ in $\mathbf{T}$ in such a way that the elements $o^{\mathbf{T}}$ ($o \in \mathcal{O}$) represent every α -class.

- For every $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -module $\mathbf{Q}$, let

$$\mathbf{h}_{\mathbf{Q}} := \sum \{ |\text{Hom}(\mathbf{M}, \mathbf{Q})| : \mathbf{M} \in \text{Mod}(\mathbf{A}, \mathbf{p}) \},$$

and let $\mathbf{h}$ be the maximum of the numbers $\mathbf{h}_{\mathbf{Q}}$ as $\mathbf{Q}$ runs over all subdirectly irreducible $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -modules. ($\mathbf{h}$ stands for ‘homomorphisms’.)

- Let $\mathbf{e}$ be the least common multiple of the exponents of the additive groups of all modules $\mathbf{M} \in \text{Mod}(\mathbf{A}, \mathbf{p})$. ($\mathbf{e}$ stands for ‘exponent’.)
- Let $\mathbf{c}_0 := \max(1 + \mathbf{p}, k - 1)$, and let $\mathbf{c} := \max(\mathbf{c}_0, \mathbf{p}\mathbf{h}(\mathbf{e} + 1))$. ($\mathbf{c}$ stands for ‘constant’.)

Note that the ring $\mathbf{R}(\mathcal{V}, \mathcal{O})$ is finite, because it consists of $\mathcal{O} \times \mathcal{O}$ matrices where each entry is determined by an element of the free algebra $\mathbf{F}$ in $\mathcal{V}$ with free generating set $\{x\} \cup \mathcal{O}$, and $\mathbf{F}$ is finite, since $\mathcal{V}$ is generated by $\mathbf{A}$ (a finite algebra). The set $\text{Mod}(\mathbf{A}, \mathbf{p})$ is also finite, since $\mathbf{A}$, $\mathbf{p}$, and $\mathcal{O}$ are finite. It follows from the main result of [12] that $|\mathbf{Q}| \leq |\mathbf{R}(\mathcal{V}, \mathcal{O})|$ holds for every subdirectly irreducible $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -module $\mathbf{Q}$. Therefore there are only finitely many subdirectly irreducible $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -modules, up to isomorphism, and all are finite, so $\mathbf{h}$ is a natural number. The finiteness of $\text{Mod}(\mathbf{A}, \mathbf{p})$ implies also that $\mathbf{e}$ is a natural number, and hence so are $\mathbf{c}_0$ and $\mathbf{c}$.

Remarks 5.1. (a) The definition of $\text{Mod}(\mathbf{A}, \mathbf{p})$ shows that $\text{Mod}(\mathbf{A}, \mathbf{p}) \neq \emptyset$ if and only if some power $\mathbf{A}^p$ of $\mathbf{A}$ with $1 \leq p \leq \mathbf{p}$ has a subalgebra that is isomorphic to a section of $\mathbf{A}$ with a nontrivial abelian congruence.

(b) If $\text{Mod}(\mathbf{A}, \mathbf{p}) = \emptyset$, then $\mathbf{p} = 1$, $\mathbf{e} = 1$, $\mathbf{h} = 0$, and $\mathbf{c} = \mathbf{c}_0 = \max(2, k - 1)$, while if $\text{Mod}(\mathbf{A}, \mathbf{p}) \neq \emptyset$, then $\mathbf{p} \geq 1$, $\mathbf{e} \geq 2$, and $\mathbf{h} \geq |\text{Mod}(\mathbf{A}, \mathbf{p})| \geq 1$, because the summands in the definition of $\mathbf{h}_{\mathbf{Q}}$ satisfy $|\text{Hom}(\mathbf{M}, \mathbf{Q})| \geq 1$ for all $\mathbf{M} \in \text{Mod}(\mathbf{A}, \mathbf{p})$ and all $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -modules $\mathbf{Q}$.

(c) In particular, for the trivial $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -module $\mathbf{Q}_0$ we have $\mathbf{h}_{\mathbf{Q}_0} = |\text{Mod}(\mathbf{A}, \mathbf{p})|$, therefore the inequality in (b) implies that $\mathbf{h} \geq \mathbf{h}_{\mathbf{Q}_0}$.

We start the proof of Theorem 1.1 by considering compatible relations B of $\mathbf{A}$ that are constructed in Theorem 3.1. We will use the same notation for these relations as in Theorem 3.1, except that the $\sim$ ’s from the notation of the congruences $\tilde{\alpha}_i$ will be omitted. So, the set of relations we will consider, and will denote by $\mathcal{R}^*(\mathbf{A})$,

consists of all compatible relations B of $\mathbf{A}$ that satisfy the following condition $(*)$ from Theorem 3.1, for some n :

- $(*)$ There exist
 - (I) subalgebras $\mathbf{B}_i \leq \mathbf{A}^{p_i}$ with $p_i \leq \mathbf{p}$ for each $i \in [n]$ such that $\mathbf{B}_i$ is isomorphic to a section of $\mathbf{A}$, and
 - (II) nontrivial abelian congruences $\alpha_i \in \text{Con}(\mathbf{B}_i)$ ($i \in [n]$) such that
 - (III) B is the universe of a subdirect product $\mathbf{B}$ of $\mathbf{B}_1, \dots, \mathbf{B}_n$, and
 - (IV) the product congruence $\alpha := \prod_{i=1}^n \alpha_i$ of $\prod_{i=1}^n \mathbf{B}_i$ restricts to $\mathbf{B}$ as a congruence $\alpha_{\mathbf{B}}$ of index $\leq i$.

We will refer to n as the $*$ -arity of B . Since $\mathbf{B} \leq \prod_{i=1}^n \mathbf{B}_i \leq \prod_{i=1}^n \mathbf{A}^{p_i}$, the arity of B , as a compatible relation of $\mathbf{A}$, is $\sum_{i=1}^n p_i$.

Lemma 5.2. *Let $B \in \mathcal{R}^*(\mathbf{A})$ be a compatible relation of $\mathbf{A}$ of $*$ -arity n , and let $\mathbf{B}_i$, α_i , $\mathbf{B}$, and α be as in $(*)$. If $n > \mathbf{h}(\mathbf{e} + 1)$ and $\mathbf{B}$ is a $\cap$ -irreducible subalgebra of its α -saturation $\mathbf{B}[\alpha]$ in $\prod_{i=1}^n \mathbf{B}_i$, then there exist*

- (1) a compatible relation $B' \in \mathcal{R}^*(\mathbf{A})$ of $\mathbf{A}$ of $*$ -arity $\leq n - \mathbf{e}$ and
- (2) subalgebras $\mathbf{D} \leq \mathbf{B}_i^{\mathbf{e}+1}$ and $\mathbf{D}' \leq \mathbf{B}_i^2$ for some $i \in [n]$

such that

- (3) $\{B_1, \dots, B_n, B', D, D'\} \models_d B$.

Proof. Let B satisfy the hypotheses of the lemma, including the assumptions that $n > \mathbf{h}(\mathbf{e} + 1)$ and $\mathbf{B}$ is a $\cap$ -irreducible subalgebra of $\mathbf{B}[\alpha]$. Since $\mathbf{A}$ has a parallelogram term, we know from Theorem 2.5 that the variety $\mathcal{V}$ it generates is congruence modular. As in Section 4, we expand the language of $\mathcal{V}$ by $\mathcal{O}$.

Let us fix interpretations $o^{\mathbf{B}}$ for each $o \in \mathcal{O}$ in $\mathbf{B}$ in such a way that the elements $o^{\mathbf{B}}$ ($o \in \mathcal{O}$) represent all $\alpha_{\mathbf{B}}$ -classes of $\mathbf{B}$; this is possible, since one of our assumptions is that $\alpha_{\mathbf{B}}$ has index $\leq i$. So, $\mathbf{B}$ becomes an $\mathcal{O}$ -algebra in $\mathcal{V}_{\mathcal{O}}$. Now we fix interpretations $o^{\mathbf{B}_i}$ for every constant symbol $o \in \mathcal{O}$ in each $\mathbf{B}_i$ ($i \in [n]$) such that $o^{\mathbf{B}} = (o^{\mathbf{B}_1}, \dots, o^{\mathbf{B}_n})$. Thus, each $\mathbf{B}_i$ will become an $\mathcal{O}$ -algebra in $\mathcal{V}_{\mathcal{O}}$. (Note that $\mathbf{B}_i$ and $\mathbf{B}_j$ might become different $\mathcal{O}$ -algebras even if $\mathbf{B}_i$ and $\mathbf{B}_j$ are the same as algebras in the original language of $\mathcal{V}$.) This choice makes sure that $\mathbf{B}$ is a subdirect $\mathcal{O}$ -subalgebra of the $\mathcal{O}$ -product $\prod_{i=1}^n \mathbf{B}_i$ of the $\mathcal{O}$ -algebras $\mathbf{B}_1, \dots, \mathbf{B}_n$. Since the elements $o^{\mathbf{B}}$ ($o \in \mathcal{O}$) represent all $\alpha_{\mathbf{B}}$ -classes of $\mathbf{B}$, it follows that for every $i \in [n]$, the elements $o^{\mathbf{B}_i}$ ($o \in \mathcal{O}$) represent all α_i -classes of $\mathbf{B}_i$. Hence $\mathbf{M}_{\mathbf{B}_i}(\alpha_i, \mathcal{O}) \in \text{Mod}(\mathbf{A}, \mathbf{p})$ for every $i \in [n]$.

To make $\mathbf{B}[\alpha]$ into an $\mathcal{O}$ -algebra in $\mathcal{V}_{\mathcal{O}}$, let $o^{\mathbf{B}[\alpha]} := o^{\mathbf{B}}$ for every $o \in \mathcal{O}$. Then $\mathbf{B}[\alpha]$ is also a subdirect $\mathcal{O}$ -subalgebra of the $\mathcal{O}$ -product $\prod_{i=1}^n \mathbf{B}_i$ of the $\mathcal{O}$ -algebras $\mathbf{B}_1, \dots, \mathbf{B}_n$, and $\mathbf{B}$ is an $\mathcal{O}$ -subalgebra of $\mathbf{B}[\alpha]$. Moreover, since every $\alpha_{\mathbf{B}[\alpha]}$ -class contains an $\alpha_{\mathbf{B}}$ -class, the elements $o^{\mathbf{B}[\alpha]} = o^{\mathbf{B}}$ ($o \in \mathcal{O}$) represent all $\alpha_{\mathbf{B}[\alpha]}$ -classes of $\mathbf{B}[\alpha]$.

Now let us consider the $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -modules $\mathbf{M}_{\mathbf{B}}(\alpha_{\mathbf{B}}, \mathcal{O})$ and $\mathbf{M}_{\mathbf{B}[\alpha]}(\alpha_{\mathbf{B}[\alpha]}, \mathcal{O})$. The fact that $\mathbf{B}$ is an $\mathcal{O}$ -subalgebra of $\mathbf{B}[\alpha]$ implies, by Theorem 4.6, that $\mathbf{M}_{\mathbf{B}}(\alpha_{\mathbf{B}}, \mathcal{O})$ is an $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -submodule of $\mathbf{M}_{\mathbf{B}[\alpha]}(\alpha_{\mathbf{B}[\alpha]}, \mathcal{O})$. Our additional assumption that $\mathbf{B}$ is a $\cap$ -irreducible subalgebra of $\mathbf{B}[\alpha]$ implies that $\mathbf{B}$ is also $\cap$ -irreducible in the lattice of $\mathcal{O}$ -subalgebras of $\mathbf{B}[\alpha]$, hence, again by Theorem 4.6, $\mathbf{M}_{\mathbf{B}}(\alpha_{\mathbf{B}}, \mathcal{O})$ is a $\cap$ -irreducible $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -submodule of $\mathbf{M}_{\mathbf{B}[\alpha]}(\alpha_{\mathbf{B}[\alpha]}, \mathcal{O})$. Recall also that by Lemma 4.8 (applied to $\mathbf{C} = \mathbf{B}[\alpha]$), the $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -modules $\mathbf{M}_{\mathbf{B}[\alpha]}(\alpha_{\mathbf{B}[\alpha]}, \mathcal{O})$ and $\prod_{i=1}^n \mathbf{M}_{\mathbf{B}_i}(\alpha_i, \mathcal{O})$ are naturally isomorphic. Therefore we can identify the modules $\mathbf{M}_{\mathbf{B}[\alpha]}(\alpha_{\mathbf{B}[\alpha]}, \mathcal{O})$ and $\prod_{i=1}^n \mathbf{M}_{\mathbf{B}_i}(\alpha_i, \mathcal{O})$ via this isomorphism, and get that $\mathbf{M}_{\mathbf{B}}(\alpha_{\mathbf{B}}, \mathcal{O})$ is a $\cap$ -irreducible $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -submodule of $\prod_{i=1}^n \mathbf{M}_{\mathbf{B}_i}(\alpha_i, \mathcal{O})$. Thus, the quotient

$$\mathbf{Q} := \left(\prod_{i=1}^n \mathbf{M}_{\mathbf{B}_i}(\alpha_i, \mathcal{O}) \right) / \mathbf{M}_{\mathbf{B}}(\alpha_{\mathbf{B}}, \mathcal{O})$$

is a subdirectly irreducible or trivial $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -module (according to whether $\mathbf{B} < \mathbf{B}[\alpha]$ or $\mathbf{B} = \mathbf{B}[\alpha]$).

Let $\varphi: \prod_{i=1}^n \mathbf{M}_{\mathbf{B}_i}(\alpha_i, \mathcal{O}) \rightarrow \mathbf{Q}$ be the natural homomorphism. For each $i \in [n]$, the mapping $\psi_i: \mathbf{M}_{\mathbf{B}_i}(\alpha_i, \mathcal{O}) \rightarrow \mathbf{Q}$ defined by $\psi_i(z) = \varphi(0, \dots, 0, z, 0, \dots, 0)$ (with z in the i -th position) for all $z \in \mathbf{M}_{\mathbf{B}_i}(\alpha_i, \mathcal{O})$ is an $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -module homomorphism, and

$$\varphi(z_1, \dots, z_n) = \sum_{i=1}^n \psi_i(z_i) \quad \text{for all } (z_1, \dots, z_n) \in \prod_{i=1}^n \mathbf{M}_{\mathbf{B}_i}(\alpha_i, \mathcal{O}).$$

Thus, for every element $(z_1, \dots, z_n)$ of $\prod_{i=1}^n \mathbf{M}_{\mathbf{B}_i}(\alpha_i, \mathcal{O})$,

$$(5.2) \quad (z_1, \dots, z_n) \in M_{\mathbf{B}}(\alpha_{\mathbf{B}}, \mathcal{O}) \quad \text{if and only if} \quad \sum_{i=1}^n \psi_i(z_i) = 0;$$

in other words, $M_{\mathbf{B}}(\alpha_{\mathbf{B}}, \mathcal{O})$ is the solution set of the equation $\sum_{i=1}^n \psi_i(z_i) = 0$ in $\prod_{i=1}^n \mathbf{M}_{\mathbf{B}_i}(\alpha_i, \mathcal{O})$.

Since the $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -module $\mathbf{Q}$ is subdirectly irreducible or trivial, and the $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -modules $\mathbf{M}_{\mathbf{B}_i}(\alpha_i, \mathcal{O})$ all belong to $\text{Mod}(\mathbf{A}, \mathbf{p})$, the definition of $\mathbf{h}$ (combined with Remarks 5.1 (c)) makes sure that there are at most $\mathbf{h}$ distinct homomorphisms among the ψ_i 's ($i \in [n]$). Therefore our assumption $n > \mathbf{h}(\mathbf{e} + 1)$ forces that at least $\mathbf{e} + 2$ of the ψ_i 's are equal. By permuting coordinates we may assume that for $i = 1, \dots, \mathbf{e} + 2$ the ψ_i 's are equal; hence for $i = 1, \dots, \mathbf{e} + 2$ the $\mathbf{M}_{\mathbf{B}_i}(\alpha_i, \mathcal{O})$'s are equal as $\mathbf{R}(\mathcal{V}, \mathcal{O})$ -modules, the $\mathbf{B}_i$'s are equal as $\mathcal{O}$ -algebras, and the α_i 's are equal. Let

$$(5.3) \quad \overline{\psi} := \psi_1 = \dots = \psi_{\mathbf{e}+2},$$

$$(5.4) \quad \overline{\mathbf{B}} := \mathbf{B}_1 = \dots = \mathbf{B}_{\mathbf{e}+2},$$

$$(5.5) \quad \overline{\alpha} := \alpha_1 = \dots = \alpha_{\mathbf{e}+2},$$

and

$$(5.6) \quad o^{\overline{\mathbf{B}}} := o^{\mathbf{B}_1} = \dots = o^{\mathbf{B}_{e+2}} \quad \text{for every } o \in \mathcal{O}.$$

From now on we will partition the coordinates of $\prod_{i=1}^n \mathbf{B}_i$ (and its subalgebras $\mathbf{B}[\alpha]$, $\mathbf{B}$, etc.) into two blocks, the first block consisting of the first $e+1$ coordinates, and the second from coordinates $e+2, \dots, n$. Accordingly, we will use the notation $\overline{\mathbf{x}} = (x_1, \dots, x_{e+1})$ for the elements of $\overline{\mathbf{B}}^{e+1} = \prod_{i=1}^{e+1} \mathbf{B}_i$ and the notation $\mathbf{x} = (x_{e+2}, \dots, x_n)$ for the elements of $\prod_{i=e+2}^n \mathbf{B}_i$. Hence, an element of $\prod_{i=1}^n \mathbf{B}_i$ (in particular, of $\mathbf{B}$ or $\mathbf{B}[\alpha]$) will be written as $(\overline{\mathbf{x}}, \mathbf{x})$. For any tuple $(\overline{\mathbf{x}}, \mathbf{x}) \in \mathbf{B}[\alpha]$ let $\mathcal{O}_{(\overline{\mathbf{x}}, \mathbf{x})}$ denote the set of all $o \in \mathcal{O}$ such that $(\overline{\mathbf{x}}, \mathbf{x}) \in o^{\mathbf{B}[\alpha]} / \alpha_{\mathbf{B}[\alpha]}$.

Claim 5.3. *The following hold for every element $(\overline{\mathbf{x}}, \mathbf{x}) \in \mathbf{B}[\alpha]$:*

(1) $\mathcal{O}_{(\overline{\mathbf{x}}, \mathbf{x})} \neq \emptyset$, and for any $o \in \mathcal{O}$,

$$\begin{aligned} o \in \mathcal{O}_{(\overline{\mathbf{x}}, \mathbf{x})} &\Leftrightarrow x_i \in o^{\mathbf{B}_i} / \alpha_i \text{ for all } i \in [n] \\ &\Leftrightarrow x_i \in o^{\overline{\mathbf{B}}} / \overline{\alpha} \text{ for all } i \in [e+1], \text{ and} \\ &\quad x_i \in o^{\mathbf{B}_i} / \alpha_i \text{ for all } i \in [n] \setminus [e+1]. \end{aligned}$$

(2) In particular, $x_1 \equiv_{\overline{\alpha}} x_2 \equiv_{\overline{\alpha}} \dots \equiv_{\overline{\alpha}} x_{e+1}$.

Proof of Claim 5.3. Let $(\overline{\mathbf{x}}, \mathbf{x}) \in \mathbf{B}[\alpha]$.

To prove (1) observe first that since the elements $o^{\mathbf{B}[\alpha]}$ ($o \in \mathcal{O}$) represent all $\alpha_{\mathbf{B}[\alpha]}$ -classes, there is an $o \in \mathcal{O}$ such that $o^{\mathbf{B}[\alpha]}$ is in the $\alpha_{\mathbf{B}[\alpha]}$ -class of $(\overline{\mathbf{x}}, \mathbf{x})$. Hence $o \in \mathcal{O}_{(\overline{\mathbf{x}}, \mathbf{x})}$, proving that $\mathcal{O}_{(\overline{\mathbf{x}}, \mathbf{x})} \neq \emptyset$. For the second statement in (1), recall that $o^{\mathbf{B}[\alpha]} = (o^{\mathbf{B}_1}, \dots, o^{\mathbf{B}_n})$ for every $o \in \mathcal{O}$, and that $\alpha_{\mathbf{B}[\alpha]}$ is the restriction of the product congruence $\alpha = \prod_{i=1}^n \alpha_i$ to $\mathbf{B}[\alpha]$. Thus, $(\overline{\mathbf{x}}, \mathbf{x}) \in o^{\mathbf{B}[\alpha]} / \alpha_{\mathbf{B}[\alpha]}$ if and only if $x_i \in o^{\mathbf{B}_i} / \alpha_i$ for all $i \in [n]$. This, together with the definition of $\mathcal{O}_{(\overline{\mathbf{x}}, \mathbf{x})}$ implies the first $\Leftrightarrow$ in the displayed statement. The second $\Leftrightarrow$ follows by (5.4)–(5.6).

For (2), we get from (1) that if $o \in \mathcal{O}_{(\overline{\mathbf{x}}, \mathbf{x})}$ then $x_1, \dots, x_{e+1} \in o^{\overline{\mathbf{B}}} / \overline{\alpha}$. Since $\mathcal{O}_{(\overline{\mathbf{x}}, \mathbf{x})} \neq \emptyset$, this implies that $x_1, \dots, x_{e+1}$ are in the same $\overline{\alpha}$ -class. $\diamond$

Claim 5.4. *The following conditions on an element $(\overline{\mathbf{x}}, \mathbf{x}) \in \mathbf{B}[\alpha]$ are equivalent:*

- (a) $(\overline{\mathbf{x}}, \mathbf{x}) \in \mathbf{B}$;
- (b) $(y, \dots, y, \mathbf{x}) \in \mathbf{B}$ (with $e+1$ occurrences of y) for the element y defined by

$$(5.7) \quad y = d^{\overline{\mathbf{B}}}(d^{\overline{\mathbf{B}}}(\dots d^{\overline{\mathbf{B}}}(d^{\overline{\mathbf{B}}}(x_1, x_{e+1}, x_2), x_{e+1}, x_3) \dots), x_{e+1}, x_e).$$

Proof of Claim 5.4. Throughout the proof, we will work with a fixed (but arbitrary) tuple $(\overline{\mathbf{x}}, \mathbf{x}) \in \mathbf{B}[\alpha]$, and y will denote the element defined in (5.7). By Claim 5.3(1), $\mathcal{O}_{(\overline{\mathbf{x}}, \mathbf{x})} \neq \emptyset$ and we have $x_i \in o^{\mathbf{B}_i} / \alpha_i$ whenever $o \in \mathcal{O}_{(\overline{\mathbf{x}}, \mathbf{x})}$ and $i \in [n]$. Consequently, for each $o \in \mathcal{O}_{(\overline{\mathbf{x}}, \mathbf{x})}$ and $i \in [n]$, the $\mathcal{O}$ -tuple $[x_i]^o$ that has x_i in position o and zeros (i.e., $o^{\mathbf{B}_i}$) in all other positions o' is an element of $\mathbf{M}_{\mathbf{B}_i}(\alpha_i, \mathcal{O})$. Similarly,

the $\mathcal{O}$ -tuple $[(\bar{\mathbf{x}}, \mathbf{x})]^o$ that has $(\bar{\mathbf{x}}, \mathbf{x}) = (x_i)_{i \in [n]}$ in position o and zeros (i.e., $o^{\mathbf{B}[\alpha]} = (o^{\mathbf{B}_1}, \dots, o^{\mathbf{B}_n})$) in all other positions o' is an element of $\mathbf{M}_{\mathbf{B}[\alpha]}(\alpha_{\mathbf{B}[\alpha]}, \mathcal{O})$. By inspecting the isomorphism that we use to identify $\mathbf{M}_{\mathbf{B}[\alpha]}(\alpha_{\mathbf{B}[\alpha]}, \mathcal{O})$ with $\prod_{i=1}^n \mathbf{M}_{\mathbf{B}_i}(\alpha_i, \mathcal{O})$ it is easy to see that this identification yields that $[(\bar{\mathbf{x}}, \mathbf{x})]^o = ([x_i]^o)_{i \in [n]}$.

Therefore, if $(\bar{\mathbf{x}}, \mathbf{x}) \in \mathbf{B}$, then $([x_i]^o)_{i \in [n]} = [(\bar{\mathbf{x}}, \mathbf{x})]^o \in M_{\mathbf{B}}(\alpha_{\mathbf{B}}, \mathcal{O})$ for all $o \in \mathcal{O}_{(\bar{\mathbf{x}}, \mathbf{x})}$, so using (5.2) we see that (a) implies the following condition:

$$(a)' \quad \sum_{i=1}^n \psi_i([x_i]^o) = 0 \text{ for all } o \in \mathcal{O}_{(\bar{\mathbf{x}}, \mathbf{x})}.$$

Conversely, assume that (a)' holds for $(\bar{\mathbf{x}}, \mathbf{x})$. Since $\mathcal{O}_{(\bar{\mathbf{x}}, \mathbf{x})} \neq \emptyset$, we can fix an $o \in \mathcal{O}_{(\bar{\mathbf{x}}, \mathbf{x})}$ and use (5.2) to conclude that $[(\bar{\mathbf{x}}, \mathbf{x})]^o = ([x_i]^o)_{i \in [n]} \in M_{\mathbf{B}}(\alpha_{\mathbf{B}}, \mathcal{O})$. By the definition of $\mathbf{M}_{\mathbf{B}}(\alpha_{\mathbf{B}}, \mathcal{O})$, this implies that $(\bar{\mathbf{x}}, \mathbf{x}) \in \mathbf{B}$. Thus, (a) $\Leftrightarrow$ (a)'.

Now let us consider the element $y \in \bar{\mathbf{B}}$. By Claim 5.3(1), for every $o \in \mathcal{O}_{(\bar{\mathbf{x}}, \mathbf{x})}$ we have that $x_1, \dots, x_{e+1} \in o^{\bar{\mathbf{B}}}/\bar{\alpha}$, so the idempotence of d implies that $y \in o^{\bar{\mathbf{B}}}/\bar{\alpha}$. Since $\mathcal{O}_{(\bar{\mathbf{x}}, \mathbf{x})} \neq \emptyset$, it follows that $(\bar{\mathbf{x}}, \mathbf{x}) \equiv_{\alpha} (y, \dots, y, \mathbf{x})$. Thus, $(y, \dots, y, \mathbf{x}) \in \mathbf{B}[\alpha]$ and the tuples $(\bar{\mathbf{x}}, \mathbf{x})$ and $(y, \dots, y, \mathbf{x})$ belong to the same $\alpha_{\mathbf{B}[\alpha]}$ -classes $o^{\mathbf{B}[\alpha]}/\alpha_{\mathbf{B}[\alpha]}$ ($o \in \mathcal{O}$). Consequently, $\mathcal{O}_{(\bar{\mathbf{x}}, \mathbf{x})} = \mathcal{O}_{(y, \dots, y, \mathbf{x})}$.

For every $o \in \mathcal{O}_{(y, \dots, y, \mathbf{x})} = \mathcal{O}_{(\bar{\mathbf{x}}, \mathbf{x})}$ the $\mathcal{O}$ -tuple $[y]^o$ that has y in position o and zeros (i.e., $o^{\bar{\mathbf{B}}}$) in all other positions o' is an element of $\mathbf{M}_{\bar{\mathbf{B}}}(\bar{\alpha}, \mathcal{O})$, and the $\mathcal{O}$ -tuple $[(y, \dots, y, \mathbf{x})]^o$ that has $(y, \dots, y, \mathbf{x}) = (y, \dots, y, x_{e+2}, \dots, x_n)$ in position o and zeros (i.e., $o^{\mathbf{B}[\alpha]} = (o^{\mathbf{B}_1}, \dots, o^{\mathbf{B}_n})$) in all other positions o' is an element of $\mathbf{M}_{\mathbf{B}[\alpha]}(\alpha_{\mathbf{B}[\alpha]}, \mathcal{O})$. As before, the identification of $\mathbf{M}_{\mathbf{B}[\alpha]}(\alpha_{\mathbf{B}[\alpha]}, \mathcal{O})$ with $\prod_{i=1}^n \mathbf{M}_{\mathbf{B}_i}(\alpha_i, \mathcal{O})$ yields that $[(y, \dots, y, \mathbf{x})]^o = ([y]^o, \dots, [y]^o, [x_{e+2}]^o, \dots, [x_n]^o)$. Hence, if we apply the equivalence of conditions (a) and (a)' to the tuple $(y, \dots, y, \mathbf{x}) \in \mathbf{B}[\alpha]$, we obtain that (b) $\Leftrightarrow$ (b)' for the condition

$$(b)' \quad \sum_{i=1}^{e+1} \psi_i([y]^o) + \sum_{i=e+2}^n \psi_i([x_i]^o) = 0 \text{ for all } o \in \mathcal{O}_{(y, \dots, y, \mathbf{x})} = \mathcal{O}_{(\bar{\mathbf{x}}, \mathbf{x})}.$$

Thus, our claim (a) $\Leftrightarrow$ (b) will follow if we prove that (a)' $\Leftrightarrow$ (b)'.

The only difference between conditions (a)' and (b)' are in the first $e+1$ summands. Therefore we will be done if we show that

$$(5.8) \quad \sum_{i=1}^{e+1} \psi_i([x_i]^o) = \sum_{i=1}^{e+1} \psi_i([y]^o) \quad \text{for every } o \in \mathcal{O}_{(\bar{\mathbf{x}}, \mathbf{x})} = \mathcal{O}_{(y, \dots, y, \mathbf{x})}.$$

So, let $o \in \mathcal{O}_{(\bar{\mathbf{x}}, \mathbf{x})} = \mathcal{O}_{(y, \dots, y, \mathbf{x})}$. Then, by Claim 5.3(1) and by our earlier argument on y , we have that $x_1, \dots, x_{e+1}, y \in o^{\bar{\mathbf{B}}}/\bar{\alpha}$. By combining the definition of y in (5.7) with Corollary 4.2 we get that $y = x_1 +_o x_2 +_o \dots +_o x_{e+1}$ holds in the abelian group $\mathbf{M}_{\bar{\mathbf{B}}}(\bar{\alpha}, o)$. Hence, by the definition of $+$ in the module $\mathbf{M}_{\bar{\mathbf{B}}}(\bar{\alpha}, \mathcal{O})$,

$$(5.9) \quad [y]^o = [x_1]^o + [x_2]^o + \dots + [x_{e+1}]^o$$

holds in $\mathbf{M}_{\overline{\mathbf{B}}}(\overline{\alpha}, \mathcal{O})$. So, using first (5.3), then the fact that $\overline{\psi}$ is a module homomorphism $\mathbf{M}_{\overline{\mathbf{B}}}(\overline{\alpha}, \mathcal{O}) \rightarrow \mathbf{Q}$, and finally (5.9), we get that

$$\sum_{i=1}^{e+1} \psi_i([x_i]^o) = \sum_{i=1}^{e+1} \overline{\psi}([x_i]^o) = \overline{\psi}\left(\sum_{i=1}^{e+1} [x_i]^o\right) = \overline{\psi}([y]^o).$$

Similarly,

$$\sum_{i=1}^{e+1} \psi_i([y]^o) = (e+1)\overline{\psi}([y]^o) = \overline{\psi}((e+1)[y]^o) = \overline{\psi}([y]^o),$$

where the last equality is true, because the definition of e and the fact that $[y]^o \in \mathbf{M}_{\overline{\mathbf{B}}}(\overline{\alpha}, \mathcal{O}) \in \text{Mod}(\mathbf{A}, \mathfrak{p})$ ensure that e is a multiple of the additive order of $[y]^o$. This establishes (5.8), and hence completes the proof of the claim. $\diamond$

Now we define the relations B' and D that we will use to prove Lemma 5.2. Let

$$B' := \{(y, \mathbf{x}) \in \overline{\mathbf{B}} \times \prod_{i=e+2}^n B_i : (y, \dots, y, \mathbf{x}) \in B\},$$

$$D := \{(y, \overline{\mathbf{x}}) \in \overline{\mathbf{B}}^{e+2} : (5.7) \text{ holds and } x_1 \equiv_{\overline{\alpha}} \dots \equiv_{\overline{\alpha}} x_{e+1}\}.$$

Claim 5.5. *B' has the following properties.*

- (1) $\overline{\mathbf{B}}$ has an $\mathcal{O}$ -subalgebra $\overline{\mathbf{B}}'$ such that B' is the universe of a subdirect $\mathcal{O}$ -subalgebra $\mathbf{B}'$ of the $\mathcal{O}$ -product $\overline{\mathbf{B}}' \times \prod_{i=e+2}^n \mathbf{B}_i$ of the $\mathcal{O}$ -algebras $\overline{\mathbf{B}}'$ and $\mathbf{B}_i$ ($e+2 \leq i \leq n$).
- (2) All the congruences $\overline{\alpha}' := \overline{\alpha}|_{\overline{\mathbf{B}}'} \in \text{Con}(\overline{\mathbf{B}}')$ and $\alpha_i \in \text{Con}(\mathbf{B}_i)$ ($e+2 \leq i \leq n$) are abelian, and all but possibly $\overline{\alpha}'$ are nontrivial.
- (3) For the product congruence $\alpha' := \overline{\alpha}' \times \prod_{i=e+2}^n \alpha_i$ of $\overline{\mathbf{B}}' \times \prod_{i=e+2}^n \mathbf{B}_i$, the elements $o^{\mathbf{B}'}$ ($o \in \mathcal{O}$) represent all $\alpha'_{\mathbf{B}'}$ -classes of $\mathbf{B}'$, therefore α' restricts to $\mathbf{B}'$ as a congruence $\alpha'_{\mathbf{B}'}$ of index $\leq i$.

Consequently, $\mathbf{B}'$ satisfies all conditions in (I)–(IV), with the possible exception that the abelian congruence $\overline{\alpha}'$ in the first factor might be trivial. Hence, if $\overline{\alpha}'$ is a nontrivial congruence of $\overline{\mathbf{B}}'$, then $B' \in \mathcal{R}^*(\mathbf{A})$ is a compatible relation of $\mathbf{A}$ of $*$ -arity $n - e$.

Proof of Claim 5.5. Notice first that B' is a compatible relation of $\mathbf{A}$, because $B' \neq \emptyset$ by Claim 5.4, and B' is definable by a primitive positive formula using the compatible relations $\overline{\mathbf{B}} = B_1$, B_i ($e+2 \leq i \leq n$), and B . Thus the definition of B' shows that B' is, in fact, the universe of a subalgebra $\mathbf{B}'$ of $\overline{\mathbf{B}} \times \prod_{i=e+2}^n \mathbf{B}_i$. Moreover, (5.6) implies that all tuples $(o^{\overline{\mathbf{B}}}, o^{\mathbf{B}_{e+2}}, \dots, o^{\mathbf{B}_n})$ ($o \in \mathcal{O}$) are in $\mathbf{B}'$, therefore by defining $o^{\mathbf{B}'} := (o^{\overline{\mathbf{B}}}, o^{\mathbf{B}_{e+2}}, \dots, o^{\mathbf{B}_n})$ for every $o \in \mathcal{O}$, $\mathbf{B}'$ becomes an $\mathcal{O}$ -subalgebra of $\overline{\mathbf{B}} \times \prod_{i=e+2}^n \mathbf{B}_i$. Claim 5.4 implies that $\mathbf{B}$ and $\mathbf{B}'$ have the same projections onto

their last $n - (\mathbf{e} + 1)$ coordinates, so letting $\overline{\mathbf{B}}'$ denote the projection of $\mathbf{B}'$ onto its first coordinate, we get an $\mathcal{O}$ -subalgebra $\overline{\mathbf{B}}'$ of $\overline{\mathbf{B}}$ such that $\mathbf{B}'$ is a subdirect $\mathcal{O}$ -subalgebra of the $\mathcal{O}$ -product $\overline{\mathbf{B}}' \times \prod_{i=\mathbf{e}+2}^n \mathbf{B}_i$ of the $\mathcal{O}$ -algebras $\overline{\mathbf{B}}'$ and $\mathbf{B}_i$ ($\mathbf{e} + 2 \leq i \leq n$). This proves (1).

Considering $\overline{\mathbf{B}}'$ as an algebra in the original language of $\mathcal{V}$, we have $\overline{\mathbf{B}}' \leq \overline{\mathbf{B}} = \mathbf{B}_1 \leq \mathbf{A}^{p_1}$ where $p_1 \leq \mathbf{p}$. Furthermore, since $\overline{\mathbf{B}}$ is isomorphic to a section of $\mathbf{A}$, so is its subalgebra $\overline{\mathbf{B}}'$. The other algebras $\mathbf{B}_{\mathbf{e}+2}, \dots, \mathbf{B}_n$ are unchanged, therefore we get that conditions (I) and (III) hold for B' .

Condition (II) for $\mathbf{B}$ implies that $\overline{\alpha}'$ is an abelian congruence of $\overline{\mathbf{B}}'$ and α_i is a nontrivial abelian congruence of $\mathbf{B}_i$ for every i ($\mathbf{e} + 2 \leq i \leq n$). This proves (2) and that (II) holds for $\mathbf{B}'$ with the possible exception that $\overline{\alpha}'$ may be trivial.

Finally, since the elements $o^{\mathbf{B}} = (o^{\overline{\mathbf{B}}}, \dots, o^{\overline{\mathbf{B}}}, o^{\mathbf{B}_{\mathbf{e}+2}}, \dots, o^{\mathbf{B}_n})$ ($o \in \mathcal{O}$) represent all $\alpha_{\mathbf{B}}$ -classes in $\mathbf{B}$, the definition of $\mathbf{B}'$ implies that the elements $o^{\mathbf{B}'} = (o^{\overline{\mathbf{B}}}, o^{\mathbf{B}_{\mathbf{e}+2}}, \dots, o^{\mathbf{B}_n})$ ($o \in \mathcal{O}$) represent all $\alpha'_{\mathbf{B}'}$ -classes in $\mathbf{B}'$. Hence $\alpha'_{\mathbf{B}'}$ has index $\leq |\mathcal{O}| = \mathbf{i}$. This proves (3) and that condition (IV) also hold for B' .

Thus we have $B' \in \mathcal{R}^*(\mathbf{A})$ if $\overline{\alpha}'$ is a nontrivial congruence of $\overline{\mathbf{B}}'$. It is clear from the construction of B' that its $*$ -arity is $n - \mathbf{e}$, completing the proof of Claim 5.5. $\diamond$

Claim 5.6. D is the universe of a subalgebra $\mathbf{D}$ of $\overline{\mathbf{B}}^{\mathbf{e}+1}$.

Proof of Claim 5.6. The following fact will be useful: property (4.3) of the difference term d for $\overline{\mathbf{B}}$ and its abelian congruence $\overline{\alpha}$ is equivalent to saying that

$$D_1 := \{(d^{\overline{\mathbf{B}}}(u, v, w), u, v, w) \in \overline{\mathbf{B}}^4 : u \equiv_{\overline{\alpha}} v \equiv_{\overline{\alpha}} w\}$$

is the universe of a subalgebra of $\overline{\mathbf{B}}^4$. Hence D_1 is a compatible relation of $\mathbf{A}$. Observing that all four coordinates of the tuples in D_1 are $\overline{\alpha}$ -related, one can easily check that D is definable by a primitive positive formula using D_1 . Clearly, $D \neq \emptyset$. Therefore it follows that D is a compatible relation of $\mathbf{A}$. The construction of D shows that D is, in fact, the universe of a subalgebra of $\overline{\mathbf{B}}^{\mathbf{e}+1} = \mathbf{B}_1^{\mathbf{e}+1}$. $\diamond$

Claim 5.7. $\{B_1, \dots, B_n, B', D\} \models_d B$.

Proof of Claim 5.7. Let

$$W := \{(y, \overline{\mathbf{x}}, \mathbf{x}) \in \overline{\mathbf{B}}^{\mathbf{e}+2} \times \prod_{i=\mathbf{e}+2}^n \mathbf{B}_i : (y, \overline{\mathbf{x}}) \in D \text{ and } (y, \mathbf{x}) \in B'\}.$$

Our goal is to show that

- (i) The projection map $W \rightarrow \text{proj}_{[n+1] \setminus \{1\}}(W)$ that omits the first coordinate of W is one-to-one, and

(ii) its image is B , that is,

$$(5.10) \quad \text{proj}_{[n+1] \setminus \{1\}}(W) = B.$$

(i)–(ii) will imply the statement of Claim 5.7 for the following reason. By (ii), W is nonempty, and it is easy to see from the definition of W that W is definable by a quantifier-free primitive positive formula, using $B_1, \dots, B_n, B'$ and D . Hence $\{B_1, \dots, B_n, B', D\} \models_d W$. By (i) and (ii), B is obtained from W by bijective projection, so $\{W\} \models_d B$. By the transitivity of $\models_d$ we get that $\{B_1, \dots, B_n, B', D\} \models_d B$, as claimed.

To prove (i) observe that the projection map $W \rightarrow \text{proj}_{[n+1] \setminus \{1\}}(W)$ is one-to-one, because for every element $(y, \bar{x}, \mathbf{x}) \in W$ we have $(y, \bar{x}) \in D$, and hence y is uniquely determined by $\bar{x}$, via (5.7).

To prove the inclusion $\supseteq$ in (5.10), let $(\bar{x}, \mathbf{x}) \in B$ and define $y \in \bar{B}$ by (5.7). Then Claim 5.4 implies that $(y, \dots, y, \mathbf{x}) \in B$, so $(y, \mathbf{x}) \in B'$. On the other hand, by Claim 5.3(2) we have that $x_1 \equiv_{\bar{\alpha}} \dots \equiv_{\bar{\alpha}} x_{e+1}$, therefore $(y, \bar{x}) \in D$. This shows that $(y, \bar{x}, \mathbf{x}) \in W$, and hence $(\bar{x}, \mathbf{x}) \in \text{proj}_{[n+1] \setminus \{1\}}(W)$.

For the inclusion $\subseteq$ in (5.10), assume that $(\bar{x}, \mathbf{x}) \in \text{proj}_{[n+1] \setminus \{1\}}(W)$. Then there exists $y \in \bar{B}$ such that $(y, \bar{x}, \mathbf{x}) \in W$. Let us fix such a y . By the definition of W it follows that $(y, \bar{x}) \in D$ and $(y, \mathbf{x}) \in B'$. The latter implies that $(y, \dots, y, \mathbf{x}) \in \mathbf{B}$, while the former implies that $x_1 \equiv_{\bar{\alpha}} \dots \equiv_{\bar{\alpha}} x_{e+1}$ and the equality in (5.7) holds for y . Since d is idempotent, we get that $x_1 \equiv_{\bar{\alpha}} \dots \equiv_{\bar{\alpha}} x_{e+1} \equiv_{\bar{\alpha}} y$, so $(\bar{x}, \mathbf{x}) \equiv_{\alpha} (y, \dots, y, \mathbf{x})$. Since $(y, \dots, y, \mathbf{x}) \in \mathbf{B}$, this shows that $(\bar{x}, \mathbf{x}) \in B[\alpha]$. Therefore Claim 5.4 applies, and we obtain that $(\bar{x}, \mathbf{x}) \in B$. $\diamond$

Now we are ready to prove the statements (1)–(3) of Lemma 5.2. We will use the notation and the conclusions of Claims 5.5–5.7. There are two cases to consider.

CASE 1: $\bar{\alpha}' \in \text{Con}(\bar{\mathbf{B}}')$ is nontrivial.

Then we know from Claim 5.5 that $B' \in \mathcal{R}^*(\mathbf{A})$ is a compatible relation of $\mathbf{A}$ of $*$ -arity $n - e$, so (1) holds. For (2), choose $\mathbf{D}'$ to be any subalgebra of $\bar{\mathbf{B}}^2$, say $\mathbf{D}' = \bar{\mathbf{B}}^2$. Since $\bar{\mathbf{B}} = \mathbf{B}_1$, Claim 5.6 proves that (2) also holds. Finally, (3) follows from Claim 5.7 (the choice of D' is irrelevant).

CASE 2: $\bar{\alpha}' \in \text{Con}(\bar{\mathbf{B}}')$ is trivial.

By Claim 5.5, $\bar{\mathbf{B}}'$ is an $\mathcal{O}$ -subalgebra of $\bar{\mathbf{B}}$, therefore $o^{\bar{\mathbf{B}}'} = o^{\bar{\mathbf{B}}}$ for all $o \in \mathcal{O}$. Since the elements $o^{\bar{\mathbf{B}}}$ ($o \in \mathcal{O}$) represent all $\bar{\alpha}$ -classes of $\bar{\mathbf{B}}$ and $\bar{\alpha}' = \bar{\alpha}|_{\bar{\mathbf{B}}'}$ is a trivial congruence of $\bar{\mathbf{B}}'$, we get that the underlying set of $\bar{\mathbf{B}}'$ is $\bar{B}' = \mathcal{O}^{\bar{\mathbf{B}}} (= \mathcal{O}^{\bar{\mathbf{B}}'})$, and $o_1^{\bar{\mathbf{B}}} = o_2^{\bar{\mathbf{B}}}$ whenever $o_1^{\bar{\mathbf{B}}} \equiv_{\bar{\alpha}} o_2^{\bar{\mathbf{B}}}$ in $\bar{\mathbf{B}}$ ($o_1, o_2 \in \mathcal{O}$). It follows that the assignment $o^{\bar{\mathbf{B}}} \mapsto o^{\bar{\mathbf{B}}}/\bar{\alpha}$ is an isomorphism $\bar{\mathbf{B}}' \rightarrow \bar{\mathbf{B}}/\bar{\alpha}$. Hence the map $\nu: \bar{\mathbf{B}} \rightarrow \bar{\mathbf{B}}'$ that assigns to every $\bar{b} \in \bar{B}$ the unique element $o^{\bar{\mathbf{B}}} \in \mathcal{O}^{\bar{\mathbf{B}}} = \bar{B}'$ such that $o^{\bar{\mathbf{B}}} \equiv_{\bar{\alpha}} \bar{b}$ is a well-defined homomorphism with kernel $\bar{\alpha}$.

Now let $\mathbf{D}'$ denote the subalgebra of $\overline{\mathbf{B}}^2$ whose universe

$$D' = \{(x, y) \in \overline{\mathbf{B}}^2 : y = \nu(x)\}$$

is the graph of ν , and let $\mathbf{B}'' = \text{proj}_{[n-e]\setminus\{1\}}(\mathbf{B}')$ be the image of $\mathbf{B}'$ under the projection homomorphism onto all coordinates except the first. Since $\overline{\mathbf{B}} = \mathbf{B}_1$, Claim 5.6 and the definition of $\mathbf{D}'$ imply that condition (2) of Lemma 5.2 holds for $\mathbf{D}$ and $\mathbf{D}'$. It follows from the conclusions of Claim 5.5 that B'' is a compatible relation of $\mathbf{A}$ that belongs to $\mathcal{R}^*(\mathbf{A})$ and has $*$ -arity $n - e - 1$. Thus, condition (1) of Lemma 5.2 holds for B'' in place of B' . It remains to establish that condition (3) of Lemma 5.2 also holds with B'' replacing B' , that is,

$$(5.11) \quad \{B_1, \dots, B_n, B'', D, D'\} \models_d B.$$

First we prove that

$$(5.12) \quad B' = \{(y, x_{e+2}, \dots, x_n) \in \overline{\mathbf{B}} \times B'' : (x_{e+2}, y) \in D'\}.$$

To verify the inclusion $\subseteq$, let $(y, \mathbf{x}) = (y, x_{e+2}, \dots, x_n)$ be an arbitrary element of $\mathbf{B}'$. Clearly, $(y, \mathbf{x}) \in \overline{\mathbf{B}} \times B''$, so we need to show that $(x_{e+2}, y) \in D'$, that is, $y = \nu(x_{e+2})$. By Claim 5.5, $\mathbf{B}'$ is a subdirect $\mathcal{O}$ -subalgebra of $\overline{\mathbf{B}}' \times \mathbf{B}_{e+2} \times \dots \times \mathbf{B}_n$, and the elements $o^{\mathbf{B}'} = (o^{\overline{\mathbf{B}}}, o^{\mathbf{B}_{e+2}}, \dots, o^{\mathbf{B}_n})$ represent all $\alpha'_{\mathbf{B}'}$ -classes of $\mathbf{B}'$. Also, recall from (5.4)–(5.6) that at the beginning of the proof of Lemma 5.2 we arranged that $\mathbf{B}_{e+2} = \overline{\mathbf{B}}$, $\alpha_{e+2} = \overline{\alpha}$, and $o^{\mathbf{B}_{e+2}} = o^{\overline{\mathbf{B}}}$ for all $o \in \mathcal{O}$. Therefore, for the given element $(y, \mathbf{x}) \in \mathbf{B}'$, there exists $o \in \mathcal{O}$ such that $o^{\mathbf{B}'} \in (y, \mathbf{x})/\alpha'$, and so $y \equiv_{\overline{\alpha}'} o^{\overline{\mathbf{B}}} = o^{\mathbf{B}_{e+2}} \equiv_{\alpha_{e+2}} x_{e+2}$. Since $\overline{\alpha}'$ is trivial and $\mathbf{B}_{e+2} = \overline{\mathbf{B}}$, $\alpha_{e+2} = \overline{\alpha}$, we get that $y = o^{\overline{\mathbf{B}}} \equiv_{\overline{\alpha}} x_{e+2} (\in \overline{\mathbf{B}})$. Thus, $y = \nu(x_{e+2})$. This completes the proof of $\subseteq$ in (5.12).

For $\supseteq$, assume that $(y, x_{e+2}, \dots, x_n) \in \overline{\mathbf{B}} \times B''$ is such that $(x_{e+2}, y) \in D'$, that is, $y = \nu(x_{e+2})$. Since $(x_{e+2}, \dots, x_n) \in B'' = \text{proj}_{[n-e]\setminus\{1\}}(B')$, there exists $y' \in \overline{\mathbf{B}}'$ such that $(y', x_{e+2}, \dots, x_n) \in B'$. The inclusion $\subseteq$ in (5.12) proved in the preceding paragraph implies that $y' = \nu(x_{e+2})$. Thus $y = y'$ and $(y, x_{e+2}, \dots, x_n) = (y', x_{e+2}, \dots, x_n) \in B'$. This completes the proof of (5.12).

By (5.12), B' is definable by a quantifier-free primitive positive formula, using $\overline{\mathbf{B}} = B_1$, B'' and D' . Hence $\{B_1, B'', D'\} \models_d B'$. It follows that

$$\{B_1, \dots, B_n, B'', D, D'\} \models_d \{B_1, \dots, B_n, B', D\}.$$

Combining this with the result of Claim 5.7, and using the transitivity of $\models_d$, we obtain the desired conclusion (5.11).

The proof of Lemma 5.2 is complete. $\square$

Corollary 5.8. *If some power $\mathbf{A}^p$ of $\mathbf{A}$ with $1 \leq p \leq \mathbf{p}$ has a subalgebra that is isomorphic to a section of $\mathbf{A}$ with a nontrivial abelian congruence, then $\mathcal{R}_{\text{ph}(e+1)}(\mathbf{A}) \models_d \mathcal{R}^*(\mathbf{A})$.*

Proof. As we noted in Remarks 5.1(b), the assumption that some $\mathbf{A}^p$ with $1 \leq p \leq \mathbf{p}$ has a subalgebra isomorphic to a section of $\mathbf{A}$ with a nontrivial abelian congruence implies that $\mathbf{h} \geq 1$ and $\mathbf{e} \geq 2$.

To prove the corollary we have to show that

$$(5.13) \quad \mathcal{R}_{\mathbf{ph}(\mathbf{e}+1)}(\mathbf{A}) \models_{\mathbf{d}} B$$

for every $B \in \mathcal{R}^*(\mathbf{A})$. We proceed by induction on the $*$ -arity n of B . We will use the same notation for the data $\mathbf{B}_i, \alpha_i$ ($i \in [n]$) and α associated to B as in (I)–(IV). If $n \leq \mathbf{h}(\mathbf{e}+1)$, then the arity of B (as a compatible relation of $\mathbf{A}$) is $\sum_{i=1}^n p_i \leq n\mathbf{p}$, therefore $B \in \mathcal{R}_{\mathbf{ph}(\mathbf{e}+1)}(\mathbf{A})$ and (5.13) is trivial. So assume that $n > \mathbf{h}(\mathbf{e}+1)$. The algebra $\mathbf{B}$ is the intersection of a family $\mathcal{I}$ of $\cap$ -irreducible subalgebras of $\mathbf{B}[\alpha]$. For every $\widehat{\mathbf{B}} \in \mathcal{I}$ we have $\mathbf{B} \leq \widehat{\mathbf{B}} \leq \mathbf{B}[\alpha]$, therefore $\widehat{\mathbf{B}}$ is a subdirect product of $\mathbf{B}_1, \dots, \mathbf{B}_n$ and $\widehat{\mathbf{B}}[\alpha] = \mathbf{B}[\alpha]$. The latter implies that the index of $\alpha_{\widehat{\mathbf{B}}}$ in $\widehat{\mathbf{B}}$ is the same as the index of $\alpha_{\mathbf{B}}$ in $\mathbf{B}$, because both are equal to the index of $\alpha_{\mathbf{B}[\alpha]}$ in $\mathbf{B}[\alpha]$. This shows that for every $\widehat{\mathbf{B}} \in \mathcal{I}$, $\widehat{B}$ is a relation in $\mathcal{R}^*(\mathbf{A})$ with $*$ -arity n . Since $\cap$ is an $\models_{\mathbf{d}}$ -construct, it suffices to prove (5.13) for the case when $\mathbf{B}$ is a $\cap$ -irreducible subalgebra of $\mathbf{B}[\alpha]$. Then, by Lemma 5.2, $\{B_1, \dots, B_n, B', D, D'\} \models_{\mathbf{d}} B$ for some $B' \in \mathcal{R}^*(\mathbf{A})$ of $*$ -arity $\leq n - \mathbf{e}$ and some compatible relations D and D' of $\mathbf{A}$ that are universes of algebras $\mathbf{D} \leq \mathbf{B}_i^{\mathbf{e}+1} \leq \mathbf{A}^{p_i(\mathbf{e}+1)}$ and $\mathbf{D}' \leq \mathbf{B}_i^2 \leq \mathbf{A}^{2p_i}$ for some i ($i \in [n]$). Now the induction hypothesis implies that $\mathcal{R}_{\mathbf{ph}(\mathbf{e}+1)}(\mathbf{A}) \models_{\mathbf{d}} B'$, because $n - \mathbf{e} < n$, while the inequalities $\mathbf{h} \geq 1$, $\mathbf{e} \geq 2$ from the first paragraph of this proof imply that $B_1, \dots, B_n, D, D' \in \mathcal{R}_{\mathbf{ph}(\mathbf{e}+1)}(\mathbf{A})$, because $p_i \leq 2p_i \leq p_i(\mathbf{e}+1) \leq \mathbf{p}(\mathbf{e}+1) \leq \mathbf{ph}(\mathbf{e}+1)$. Hence

$$\mathcal{R}_{\mathbf{ph}(\mathbf{e}+1)}(\mathbf{A}) \models_{\mathbf{d}} \{B_1, \dots, B_n, B', D, D'\} \models_{\mathbf{d}} B,$$

so (5.13) follows by the transitivity of $\models_{\mathbf{d}}$. $\square$

Now we are ready to prove Theorem 1.1.

Proof of Theorem 1.1. Assume that $\mathbf{A}$ satisfies the hypotheses of Theorem 1.1, that is, in addition to our global assumptions in this section that $\mathbf{A}$ is a finite algebra with a k -parallelogram term, $\mathbf{A}$ also satisfies the split centralizer condition. (For the definition of the split centralizer condition, see the Introduction.) Our aim is to prove that

$$(5.14) \quad \mathcal{R}_{\mathbf{c}}(\mathbf{A}) \models_{\mathbf{d}} \mathcal{R}(\mathbf{A}).$$

In addition to the notation $\mathcal{R}^*(\mathbf{A})$ introduced before Lemma 5.2, we will write $\mathcal{R}_{\text{crit}}(\mathbf{A})$ for the set of all critical relations of $\mathbf{A}$. (Critical relations are defined in Section 2, subsection 2.2.)

First we will argue that

$$(5.15) \quad \mathcal{R}_{\text{crit}}(\mathbf{A}) \models_{\mathbf{d}} \mathcal{R}(\mathbf{A}).$$

Every compatible relation of $\mathbf{A}$ is an intersection of $\cap$ -irreducible compatible relations. Furthermore, if a $\cap$ -irreducible compatible relation of $\mathbf{A}$ is not critical (i.e., not directly indecomposable), then up to a permutation of coordinates, it has the form $\rho \times A^\ell$, so it follows that ρ is a critical relation of $\mathbf{A}$; A itself is clearly a critical relation of $\mathbf{A}$. Therefore every compatible relation of $\mathbf{A}$ can be obtained from critical relations of $\mathbf{A}$ by product and intersection. Since product and intersection are $\models_d$ -constructs, (5.15) follows.

Theorem 3.1 implies that

$$(5.16) \quad \mathcal{R}_{c_0}(\mathbf{A}) \cup \mathcal{R}^*(\mathbf{A}) \models_d \mathcal{R}_{\text{crit}}(\mathbf{A}),$$

as we will show now. Let C be a critical relation of $\mathbf{A}$ of arity n . It is clear that $\mathcal{R}_{c_0}(\mathbf{A}) \cup \mathcal{R}^*(\mathbf{A}) \models_d C$ if $n \leq c_0$ (i.e., if $C \in \mathcal{R}_{c_0}(\mathbf{A})$), so let us assume that $n > c_0$. It follows from the definition of c_0 that $n \geq \max(3, k)$. Since $\mathbf{A}$ has a k -parallelogram term and satisfies the split centralizer condition, all hypotheses of Theorem 3.1 are satisfied. Hence the theorem yields the existence of a compatible relation B of $\mathbf{A}$ such that $B \in \mathcal{R}^*(\mathbf{A})$ and $\mathcal{R}_{1+p}(\mathbf{A}) \cup \{B\} \models_d C$. Since $1 + p \leq c_0$, this implies that $\mathcal{R}_{c_0}(\mathbf{A}) \cup \mathcal{R}^*(\mathbf{A}) \models_d C$, and proves (5.16).

If no power $\mathbf{A}^p$ ($1 \leq p \leq p$) of $\mathbf{A}$ has a subalgebra that is isomorphic to a section of $\mathbf{A}$ with a nontrivial abelian congruence, then $\mathcal{R}^*(\mathbf{A}) = \emptyset$ and $\text{Mod}(\mathbf{A}, p) = \emptyset$. As we saw in Remarks 5.1(b), the latter implies that $h = 0$ and hence $c = c_0$. Thus, in this case (5.15) and (5.16) combine to show that

$$\mathcal{R}_c(\mathbf{A}) = \mathcal{R}_{c_0}(\mathbf{A}) \stackrel{(5.16)}{\models_d} \mathcal{R}_{\text{crit}}(\mathbf{A}) \stackrel{(5.15)}{\models_d} \mathcal{R}(\mathbf{A}).$$

Hence (5.14) follows by the transitivity of $\models_d$.

In the opposite case, when some $\mathbf{A}^p$ ($1 \leq p \leq p$) has a subalgebra isomorphic to a section of $\mathbf{A}$ with a nontrivial abelian congruence, Corollary 5.8 shows that

$$(5.17) \quad \mathcal{R}_{\text{ph}(e+1)}(\mathbf{A}) \models_d \mathcal{R}^*(\mathbf{A}).$$

Since $c = \max(c_0, \text{ph}(e+1))$, we have that $\mathcal{R}_{c_0}(\mathbf{A}) \cup \mathcal{R}_{\text{ph}(e+1)}(\mathbf{A}) \subseteq \mathcal{R}_c(\mathbf{A})$, so (5.15), (5.16), and (5.17) together imply that

$$\mathcal{R}_c(\mathbf{A}) \stackrel{(5.17)}{\models_d} \mathcal{R}_c(\mathbf{A}) \cup \mathcal{R}^*(\mathbf{A}) \stackrel{(5.16)}{\models_d} \mathcal{R}_{\text{crit}}(\mathbf{A}) \stackrel{(5.15)}{\models_d} \mathcal{R}(\mathbf{A}).$$

By the transitivity of $\models_d$ this shows that (5.14) holds.

This proves (5.14) in all cases. In view of Theorem 2.4, (5.14) is sufficient to conclude that $\mathbf{A}$ is dualizable. $\square$

6. APPLICATIONS

In this section we apply the main theorem of the paper to establish dualizability within some well known classes of algebras. Some of these results were known before.

In the first part of the section we identify some conditions on a variety $\mathcal{V}$ which guarantee that every finite member of $\mathcal{V}$ is dualizable. We end the section by proving that if $\mathbf{A}$ is a finite algebra with a parallelogram term and $\mathbf{SP}(\mathbf{A})$ is a variety, then $\mathbf{A}$ is dualizable (although some of the other algebras in $\mathbf{SP}(\mathbf{A})$ need not be dualizable).

We will often apply Theorem 1.1 to prove dualizability simultaneously for all members of a class $\mathcal{C}$ of finite algebras with parallelogram terms, where $\mathcal{C}$ is closed under taking subalgebras. In such cases, to get the desired conclusion, it will be enough to check that for every $\mathbf{A} \in \mathcal{C}$, each relevant triple (δ, θ, ν) of $\mathbf{A}$ is split by a triple (α, β, κ) relative to $\mathbf{SP}(\mathbf{A})$. Indeed, if every $\mathbf{A} \in \mathcal{C}$ has this property, then for every $\mathbf{A} \in \mathcal{C}$ and $\mathbf{B} \leq \mathbf{A}$ we have that $\mathbf{B} \in \mathcal{C}$, so every relevant triple of $\mathbf{B}$ is split by a triple relative to $\mathbf{SP}(\mathbf{B})$, and therefore relative to $\mathbf{SP}(\mathbf{A})$ as well. This shows that every $\mathbf{A} \in \mathcal{C}$ satisfies the split centralizer condition, which implies by Theorem 1.1 that every $\mathbf{A} \in \mathcal{C}$ is dualizable.

We will refer to some commutator identities by the number assigned to them in [10, Chapter 8]:

$$\begin{aligned} \text{(C1)} \quad & [x \wedge y, y] = x \wedge [y, y], \\ \text{(C3)} \quad & [x, y] = x \wedge y. \\ \text{(C8)} \quad & [1, x] = x. \end{aligned}$$

An algebra satisfying a given (Ci), $i \in \{1, 3, 8\}$, may be called a *(Ci)-algebra*. It is known that if $\mathcal{V}$ is congruence modular, then the subclass of (Ci)-algebras in $\mathcal{V}$ is closed under the formation of finite subdirect products and quotient algebras. (See [10, Chapter 8].)

We have met (C1) before: any residually small congruence modular variety consists of (C1)-algebras, and conversely any congruence modular variety generated by a finite algebra whose subalgebras are all (C1) is residually small.

(C3)-algebras are also called *neutral*. From [10, Chapter 8] we know that an algebra is neutral if and only if it has no nontrivial abelian congruence intervals. Therefore, we will call an interval in a congruence lattice *neutral* if it has no nontrivial abelian subintervals.

Lemma 6.1. *Let $\mathcal{V}$ be a congruence modular variety in which every finite subdirectly irreducible algebra is either solvable or is a (C8)-algebra. If $\mathbf{A} \in \mathcal{V}$ is a finite algebra, then*

- (1) *$\mathbf{A}$ has a unique pair (σ, ρ) of complementary factor congruences such that $\mathbf{A} \cong \mathbf{A}/\sigma \times \mathbf{A}/\rho$, $\mathbf{A}/\sigma$ is solvable and $\mathbf{A}/\rho$ is a (C8)-algebra; moreover*
- (2) *every congruence χ on $\mathbf{A}$ is a product congruence relative to the factorization $\mathbf{A} \cong \mathbf{A}/\sigma \times \mathbf{A}/\rho$, meaning that $\chi = (\chi \vee \sigma) \wedge (\chi \vee \rho)$.*

Proof. To prove (1) recall that the classes of solvable algebras and (C8)-algebras in $\mathcal{V}$ are closed under finite subdirect products and quotients. Therefore, if $\mathbf{A} \in \mathcal{V}$ is any

finite algebra, then it has a least congruence σ such that $\mathbf{A}/\sigma$ is solvable and also a least congruence ρ such that $\mathbf{A}/\rho$ is a (C8)-algebra. The congruence σ is contained in the kernel of any homomorphism of $\mathbf{A}$ onto a solvable subdirectly irreducible algebra and the congruence ρ is contained in the kernel of any homomorphism of $\mathbf{A}$ onto a (C8) subdirectly irreducible algebra. Hence $\sigma \wedge \rho$ is contained in the kernel of any homomorphism of $\mathbf{A}$ onto a subdirectly irreducible algebra, implying that $\sigma \wedge \rho = 0$.

The algebra $\mathbf{A}/(\sigma \vee \rho)$ is a quotient of the solvable algebra $\mathbf{A}/\sigma$ and is also a quotient of the (C8)-algebra $\mathbf{A}/\rho$, so it is both solvable and (C8). This forces it to be trivial, and therefore $\sigma \vee \rho = 1$.

Congruences σ and ρ permute, since σ is cosolvable and cosolvable congruences permute with all congruences by [10, Theorem 6.2]. This completes the proof that (σ, ρ) is a pair of complementary factor congruences, and the argument shows that $\mathbf{A}/\sigma$ is solvable and $\mathbf{A}/\rho$ is (C8).

If (σ', ρ') were a second pair of complementary factor congruences such that $\mathbf{A} \cong \mathbf{A}/\sigma' \times \mathbf{A}/\rho'$, $\mathbf{A}/\sigma'$ is solvable and $\mathbf{A}/\rho'$ is a (C8)-algebra, then the solvability of $\mathbf{A}/\sigma'$ implies that $\sigma' \supseteq \sigma$, and a similar argument shows that $\rho' \supseteq \rho$. But if (σ, ρ) and (σ', ρ') are pairs of complementary factor congruences where $\sigma \subseteq \sigma'$ and $\rho \subseteq \rho'$, then $\sigma = \sigma'$ and $\rho = \rho'$.

For (2) notice that the set of product congruences on $\mathbf{A} \cong \mathbf{A}/\sigma \times \mathbf{A}/\rho$ is closed under meet, so to prove claim (2) it suffices to show that the meet irreducible congruences on $\mathbf{A}$ are product congruences. Each one has been shown to be above σ or ρ , so is in fact a product congruence. $\square$

Under the assumptions of Lemma 6.1 a congruence χ of $\mathbf{A}$ is central (i.e., satisfies $[1, \chi] = 0$) if and only if the congruence $\bar{\chi} := (\chi \vee \sigma)/\sigma$ of $\mathbf{A}/\sigma$ is central and $\chi \leq \rho$. This can be verified as follows. By Lemma 6.1(2), χ is a product congruence relative to the factorization $\mathbf{A} \cong \mathbf{A}/\sigma \times \mathbf{A}/\rho$, therefore it follows that χ is a central congruence of $\mathbf{A}$ if and only if $\bar{\chi} := (\chi \vee \sigma)/\sigma$ is a central congruence of $\mathbf{A}/\sigma$ and $(\chi \vee \rho)/\rho$ is a central congruence of $\mathbf{A}/\rho$. But $\mathbf{A}/\rho$ is a (C8)-algebra, therefore $(\chi \vee \rho)/\rho$ is central if and only if it is the trivial congruence of $\mathbf{A}/\rho$, that is, $\chi \leq \rho$.

The statement in the preceding paragraph implies that if ζ is the center of $\mathbf{A}$, then $\bar{\zeta} = (\zeta \vee \sigma)/\sigma$ is the center of $\mathbf{A}/\sigma$ and $\zeta \leq \rho$. Consequently, if $0 =: \zeta_0 \leq \zeta =: \zeta_1 \leq \zeta_2 \leq \dots$ is the ascending central series of $\mathbf{A}$, then $0 = \bar{\zeta}_0 \leq \bar{\zeta} = \bar{\zeta}_1 \leq \bar{\zeta}_2 \leq \dots$ is the ascending central series of $\mathbf{A}/\sigma$ and $\zeta_i \leq \rho$ for all i .

Corollary 6.2. *If $\mathbf{A}$ is a finite algebra in a congruence modular variety $\mathcal{V}$ such that every finite subdirectly irreducible algebra in $\mathcal{V}$ is either nilpotent or (C8), then in the factorization $\mathbf{A} \cong \mathbf{A}/\sigma \times \mathbf{A}/\rho$ in Lemma 6.1 the first factor $\mathbf{A}/\sigma$ is nilpotent, and σ and ρ are the final congruences in the descending and ascending central series of $\mathbf{A}$, respectively.*

Proof. By the construction in Lemma 6.1, σ is the least congruence such that $\mathbf{A}/\sigma$ is solvable, so $\mathbf{A}/\sigma$ is a subdirect product of solvable subdirectly irreducible algebras in

$\mathcal{V}$. Since every solvable subdirectly irreducible algebra in $\mathcal{V}$ is nilpotent, we get that $\mathbf{A}/\sigma$ is nilpotent, and σ is the least congruence such that $\mathbf{A}/\sigma$ is nilpotent. Hence σ is the final congruence in the descending central series of $\mathbf{A}$.

To prove our claim on ρ , let $0 = \zeta_0 \leq \zeta = \zeta_1 \leq \zeta_2 \leq \dots$ be the ascending central series of $\mathbf{A}$. As we saw earlier, this implies that $0 = \bar{\zeta}_0 \leq \bar{\zeta} = \bar{\zeta}_1 \leq \bar{\zeta}_2 \leq \dots$ is the ascending central series of $\mathbf{A}/\sigma$ and $\zeta_i \leq \rho$ for all i . Since $\mathbf{A}/\sigma$ is nilpotent, $\bar{\zeta}_c = 1$ for some c , so $\zeta_c \vee \sigma = 1$ and $\zeta_c \leq \rho$. This implies that $\rho = 1 \wedge \rho = (\zeta_c \vee \sigma) \wedge \rho \stackrel{\text{mod}}{=} \zeta_c \vee (\sigma \wedge \rho) = \zeta_c \vee 0 = \zeta_c$. Moreover, $\rho = \zeta_c \leq \zeta_i \leq \rho$ for every $i \geq c$, therefore $\rho = \zeta_c$ is the final congruence in the ascending central series of $\mathbf{A}$. $\square$

We will only use the following special case.

Corollary 6.3. *If $\mathbf{A}$ is a finite algebra in a congruence modular variety $\mathcal{V}$ such that every finite subdirectly irreducible algebra in $\mathcal{V}$ is either abelian or (C8), then in the factorization $\mathbf{A} \cong \mathbf{A}/\sigma \times \mathbf{A}/\rho$ in Lemma 6.1 the first factor $\mathbf{A}/\sigma$ is abelian, $\sigma = [1, 1]$ is the derived congruence, and $\rho = \zeta$ is the center.*

Proof. Restricting the argument in the proof of Corollary 6.2 to the case when all finite nilpotent subdirectly irreducible algebras are abelian yields that $\mathbf{A}/\sigma$ is abelian and σ is the least congruence such that $\mathbf{A}$ is abelian, so $\sigma = [1, 1]$. Moreover, the center of $\mathbf{A}/\sigma$ is $1 = \bar{\zeta} = \bar{\zeta}_1$, showing that we can use $c = 1$ in the argument. Therefore $\rho = \zeta_1 = \zeta$ is the center of $\mathbf{A}$. $\square$

We extend the “neutral” terminology by calling a subdirectly irreducible algebra $\mathbf{A}$ *almost neutral* if it is nonabelian and fails (C3) ($[x, y] = x \wedge y$) in exactly one way: $[\mu, \mu] = 0$, where μ is the monolith of $\mathbf{A}$. Equivalently, a subdirectly irreducible algebra $\mathbf{A}$ with monolith μ is almost neutral if it is nonabelian and

- (i) $(0 : \mu) = \mu$, and
- (ii) the congruence interval $[[\mu, 1]]$ is neutral.

Theorem 6.4. *Let $\mathcal{V}$ be a variety with a parallelogram term. Assume that every finite subdirectly irreducible algebra in $\mathcal{V}$ is abelian, neutral or almost neutral. Then every finite algebra in $\mathcal{V}$ is dualizable.*

Proof. Any neutral or almost neutral subdirectly irreducible algebra is a (C8)-algebra, hence the finite subdirectly irreducible algebras in $\mathcal{V}$ are abelian or (C8). From Corollary 6.3 we have that any finite algebra $\mathbf{A} \in \mathcal{V}$ has the properties that

- (a) $[1, 1]$ and ζ are complementary factor congruences,
- (b) $\mathbf{A}/[1, 1]$ is abelian and $\mathbf{A}/\zeta$ is a (C8)-algebra, and
- (c) every meet irreducible congruence on $\mathbf{A}$ lies above $[1, 1]$ or ζ .

Our task is to verify that for every relevant triple (δ, θ, ν) of $\mathbf{A}$, where δ is a meet irreducible congruence, $\delta \prec \theta$, $\nu = (\delta : \theta)$, and θ/δ is abelian, there exists a triple

(α, β, κ) which splits (δ, θ, ν) relative to $\mathcal{Q} := \mathbf{SP}(\mathbf{A})$. The conditions (i)–(v) that define what this means for (α, β, κ) are listed in the Introduction.

CASE 1: $[1, 1] \leq \delta$.

We have $[1, \theta] \leq [1, 1] \leq \delta$, proving that $1 \leq (\delta : \theta)$. Hence $1 = (\delta : \theta) = \nu$. If we choose $(\alpha, \beta, \kappa) = (\zeta, [1, 1], 0)$, then we have

- (i) $\kappa = 0$ is a $\mathcal{Q}$ -congruence,
- (ii) $\beta = [1, 1] \leq \delta$,
- (iii) $\alpha \wedge \beta = \zeta \wedge [1, 1] = 0 = \kappa$,
- (iv) $\alpha \vee \beta = \zeta \vee [1, 1] = 1 = \nu$, and
- (v) $[\alpha, \alpha] = [\zeta, \zeta] = 0 \leq \kappa$.

Hence the conditions required for (α, β, κ) are met.

CASE 2: $\zeta \leq \delta$.

In this case, $\mathbf{A}/\delta$ is a nonabelian subdirectly irreducible algebra. Since (δ, θ, ν) is relevant it must be that the monolith $\mu = \theta/\delta$ of $\mathbf{A}/\delta$ is abelian. $\mathbf{A}/\delta$ must be almost neutral, hence

$$\theta/\delta \leq \nu/\delta = (0 : \mu) = \mu = \theta/\delta,$$

and therefore $\nu = \theta$.

In this case it is our aim to show that there is a congruence γ covering ζ such that $\gamma \not\leq \delta$. For such a congruence we have a perspectivity $\llbracket \zeta, \gamma \rrbracket \nearrow \llbracket \delta, \theta \rrbracket$. In this situation $(\alpha, \beta, \kappa) = (\gamma \wedge [1, 1], \delta, 0)$ is a splitting triple for (δ, θ, ν) , since

- (i) $\kappa = 0$ is a $\mathcal{Q}$ -congruence,
- (ii) $\beta = \delta$ (we need only $\beta \leq \delta$ here),
- (iii) $\beta \wedge \alpha = \delta \wedge \gamma \wedge [1, 1] = \zeta \wedge [1, 1] = 0 = \kappa$,
- (iv) $\beta \vee \alpha = \delta \vee (\gamma \wedge [1, 1]) = \delta \vee \zeta \vee (\gamma \wedge [1, 1]) \stackrel{\text{mod}}{=} \delta \vee (\gamma \wedge (\zeta \vee [1, 1])) = \delta \vee \gamma = \theta = \nu$,
and
- (v) $[\alpha, \alpha] \leq \kappa$. (There are perspectivities $\llbracket \delta, \theta \rrbracket \searrow \llbracket \zeta, \gamma \rrbracket \searrow \llbracket 0, \alpha \rrbracket = \llbracket \kappa, \alpha \rrbracket$ and the first is abelian, so the last is.)

To reiterate the conclusion just drawn, it suffices to show that $\mathbf{A}$ has a congruence γ covering ζ such that $\gamma \not\leq \delta$. Since δ is also above ζ , we may work modulo ζ and henceforth assume that $\mathbf{A}$ is a (C8)-algebra. In this situation, every subdirectly irreducible quotient of $\mathbf{A}$ is neutral or almost neutral. So, we will be done if we prove the following claim.

Claim 6.5. *If $\mathbf{A}$ is a finite (C8)-algebra in $\mathcal{V}$, and (δ, θ, ν) is a relevant triple of $\mathbf{A}$, then $\mathbf{A}$ has an atomic congruence γ such that $\gamma \not\leq \delta$.*

Proof of Claim 6.5. Let δ_i , $i = 1, \dots, n$, be the set of meet irreducible congruences of $\mathbf{A}$, and for each i let θ_i be the upper cover of δ_i . For each i choose $\nabla_i \in \{\delta_i, \theta_i\}$

according to

$$\nabla_i = \begin{cases} \delta_i & \text{if } \mathbf{A}/\delta_i \text{ is neutral;} \\ \theta_i & \text{if } \mathbf{A}/\delta_i \text{ is almost neutral.} \end{cases}$$

Equivalently, $\nabla_i \in \{\delta_i, \theta_i\}$ is chosen as small as possible so that the interval $[\nabla_i, 1]$ is neutral.

Let $\nabla = \bigcap_{i=1}^n \nabla_i$. Since each $\mathbf{A}/\nabla_i$ is neutral and the class of neutral algebras in $\mathcal{V}$ is closed under finite subdirect products we get that the interval $[\nabla, 1]$ is neutral. $\mathbf{A}$ itself is not a neutral algebra, since it has a relevant triple (δ, θ, ν) (and therefore an abelian congruence interval $[\delta, \theta]$), so $0 < \nabla$.

For each i , define $\Delta_i = \delta_i \wedge \nabla$. If $\delta_i = \nabla_i$ ($\geq \nabla$) for some i , then $\Delta_i = \nabla$. In the alternative case where $\theta_i = \nabla_i$, the interval $[\Delta_i, 1]$ contains the abelian prime quotient $[\delta_i, \theta_i]$, and $[\nabla, 1]$ contains no such prime quotient, so $\Delta_i \neq \nabla$. Hence $\nabla \not\leq \delta_i$, so there is a perspectivity $[\delta_i, \theta_i] \searrow [\Delta_i, \nabla]$. Thus, for each i , $\Delta_i = \nabla$ if $\mathbf{A}/\delta_i$ is neutral, and Δ_i is a lower cover of ∇ for which the interval $[\Delta_i, \nabla]$ is abelian if $\mathbf{A}/\delta_i$ is almost neutral.

The intersection $\bigcap_{i=1}^n \Delta_i$ equals $\bigcap_{i=1}^n \delta_i = 0$. Since the former is a meet of lower covers of ∇ , and the interval $[0, \nabla]$ is modular, this interval is complemented. Thus, for every i where $\Delta_i < \nabla$ there is an atom γ_i below ∇ such that $[0, \gamma_i] \nearrow [\Delta_i, \nabla]$. Choose the value of i where δ_i is the first congruence in the relevant triple (δ, θ, ν) . Since θ/δ is abelian, $\mathbf{A}/\delta_i = \mathbf{A}/\delta$ is almost neutral. Therefore $[0, \gamma_i] \nearrow [\Delta_i, \nabla] \nearrow [\delta_i, \theta_i] = [\delta, \theta]$. For this i we have found an atom $\gamma = \gamma_i$ satisfying $\gamma \not\leq \delta$. $\diamond$

This completes the proof of the theorem. $\square$

Corollary 6.6. *Let $\mathcal{V}$ be a variety with a parallelogram term. If every finite subdirectly irreducible algebra in $\mathcal{V}$ is abelian or neutral, then every finite algebra in $\mathcal{V}$ is dualizable.*

Proof. This is Theorem 6.4 restricted to the situation where $\mathcal{V}$ has no finite almost neutral subdirectly irreducible algebras. $\square$

Corollary 6.7 (See [7]). *Any finite algebra with a near unanimity term is dualizable.*

Proof. A near unanimity term is a parallelogram term, and any algebra having such a term is neutral. Hence this corollary is a further restriction of Theorem 6.4. (In fact, it is exactly the restriction of Theorem 6.4 to the situation where $\mathcal{V}$ has no finite abelian or almost neutral subdirectly irreducible algebras.) $\square$

Corollary 6.8. *Any finite algebra in a directly representable variety is dualizable.*

Proof. This is a corollary to Corollary 6.6. To see this, recall that (i) a finite algebra in a directly representable variety has a Maltsev term (see [15, Theorem 5.11]), (ii) a Maltsev term is a parallelogram term, and (iii) the finite nonabelian subdirectly irreducible algebras in a directly representable variety are simple, hence neutral (see [15, Theorem 5.11]). $\square$

Corollary 6.9 (Cf. [6]). *Any finite algebra in a variety generated by a paraprimal algebra is dualizable.*

Proof. This follows from Corollary 6.8 and the fact that a variety generated by a paraprimal algebra is directly representable (see [4, Theorem 1.6]). $\square$

Corollary 6.10. *Any finite affine algebra is dualizable. In particular, any finite module is dualizable.*

Proof. This is another restriction of Theorem 6.4. This time we are restricting to the case where all finite subdirectly irreducible algebras in $\mathcal{V}$ are abelian. $\square$

Modules and affine algebras for which dualizability was known before include finite abelian groups (using the restriction of Pontryagin duality, see [2, Chapter 4]), finite affine spaces (see [18]), and finite algebras in a variety generated by a finite simple affine algebra (see [6]). All these special cases of Corollary 6.10 are covered also by Corollary 6.8. However, Corollary 6.10 is not a consequence of Corollary 6.8, because not every finite module lies in a directly representable variety. (A finite faithful R -module generates a directly representable variety if and only if R is of finite representation type.)

Corollary 6.11. *Let $\mathbb{K}$ be a commutative unital ring. Let $\mathcal{V}$ be a residually small variety of $\mathbb{K}$ -algebras (commutative or not, unital or not). Any finite algebra in $\mathcal{V}$ is dualizable.*

Proof. It follows from Theorems 3.1 and 3.2 in [14] that for every nonabelian subdirectly irreducible $\mathbb{K}$ -algebra S in a residually small variety the ring reduct of S is also subdirectly irreducible and lies in a residually small variety. The possible structure of a subdirectly irreducible ring in a residually small variety is described in Section 7 of [14]. These results imply that every nonabelian subdirectly irreducible algebra $S \in \mathcal{V}$ is either simple or else (i) has abelian monolith equal to the radical, $J = \text{rad}(S)$, and (ii) has S/J isomorphic to a field or a product of two fields. This is enough to guarantee that every subdirectly irreducible algebra in $\mathcal{V}$ is abelian, neutral or almost neutral. So, our statement follows from Theorem 6.4. $\square$

Corollary 6.12. *Any finite ring (commutative or not, unital or not) that generates a residually small variety is dualizable.*

Proof. This is Corollary 6.11 restricted to the case when $\mathbb{K} = \mathbb{Z}$. $\square$

The special case of Corollary 6.12 when the ring is assumed to be commutative and unital was proved in [3]. For a commutative ring R with unit, the condition that R generates a residually small variety is equivalent to the condition that its radical, $J = \text{rad}(R)$, is abelian (i.e., $J^2 = 0$).

For a finite group G , the condition that G generates a residually small variety is equivalent to the condition that the Sylow subgroups of G are abelian. Now we

explain how to derive from Theorem 1.1 the result that any finite group with abelian Sylow subgroups is dualizable. (This does not follow from Theorem 6.4.)

The crucial result is the following.

Theorem 6.13 (From [17, Chapter 2]). *If G is a finite group with abelian Sylow subgroups and (δ, θ, ν) is a relevant triple of G , then there is an endomorphism $\varepsilon: G \rightarrow G$ such that*

$$[\nu, \nu] \leq \ker(\varepsilon) \leq \delta.$$

Corollary 6.14 (From [17]). *Any finite group with abelian Sylow subgroups is dualizable.*

Proof. Let G be a finite group with abelian Sylow subgroups. We have to show that every relevant triple (δ, θ, ν) of G is split by a triple (α, β, κ) relative to $\mathcal{Q} = \mathbf{SP}(G)$. Choose $(\alpha, \beta, \kappa) = (\nu, \ker(\varepsilon), \ker(\varepsilon))$ where ε is the endomorphism from Theorem 6.13. Then

- (i) κ is a $\mathcal{Q}$ -congruence, since κ is the kernel of an endomorphism of G .
- (ii) $\beta = \ker(\varepsilon) \leq \delta$,
- (iii) $\alpha \wedge \beta = \nu \wedge \ker(\varepsilon) = \ker(\varepsilon) = \kappa$,
- (iv) $\alpha \vee \beta = \nu \vee \ker(\varepsilon) = \nu$, and
- (v) $[\alpha, \alpha] = [\nu, \nu] \leq \ker(\varepsilon) = \kappa$.

Hence the conditions required for (α, β, κ) are met. $\square$

Example 6.15. Paweł Idziak proved in [11] that the algebra $\mathbf{A}$ obtained from the six-element symmetric group S_3 by adjoining all six nullary operations is not a dualizable algebra. This is in contrast to Corollary 6.14, which establishes that S_3 without the additional constants *is* dualizable. It is worth pointing out how Idziak's example fails the conditions of Theorem 1.1. The algebra $\mathbf{A}$ has three congruences, $0 < \delta < 1$, where δ is the congruence on S_3 corresponding to the alternating group. The triple $(\delta, \theta, \nu) = (\delta, 1, 1)$ is relevant. For $\mathcal{Q} = \mathbf{SP}(\mathbf{A})$, the only $\mathcal{Q}$ -congruences on $\mathbf{A}$ are 0 and 1, and $1 \not\leq \delta$, so if (α, β, κ) is a splitting triple for (δ, θ, ν) then

- (i) κ must equal 0.

To establish dualizability our theorem also requires that

- (ii) $\beta \leq \delta$,
- (iii) $\alpha \wedge \beta = \kappa (= 0)$,
- (iv) $\alpha \vee \beta = \nu (= 1)$, and
- (v) $[\alpha, \alpha] \leq \kappa (= 0)$.

But δ is the largest abelian congruence on $\mathbf{A}$, so $\alpha \leq \delta$. This and property (ii) yield that $\alpha \vee \beta \leq \delta < 1$, contrary to (iv).

If we had not added the additional nullary operations, then the congruence δ would be another $\mathcal{Q}$ -congruence and in this situation $(\alpha, \beta, \kappa) = (1, \delta, \delta)$ would be a splitting triple for $(\delta, 1, 1)$.

We close with a different kind of application of Theorem 1.1.

Theorem 6.16. *If $\mathbf{A}$ is a finite algebra with a parallelogram term and $\mathbf{SP}(\mathbf{A})$ is a variety, then $\mathbf{A}$ is dualizable.*

Proof. If $\mathcal{Q} := \mathbf{SP}(\mathbf{A})$ is a variety, then it must be residually small, hence satisfies (C1) (by Theorems 2.1 and 2.5). Every congruence on every algebra in $\mathcal{Q}$ is a $\mathcal{Q}$ -congruence, so if (δ, θ, ν) is relevant, then it is split by (ν, δ, δ) , because (C1) implies that $[\nu, \nu] \leq \delta$. $\square$

Here it is worth pointing out that if $\mathcal{V}$ is a variety with only finitely many subdirectly irreducible algebras, each finite and having a 1-element subalgebra, then the product $\mathbf{A}$ of all these subdirectly irreducible algebras is a finite algebra with the property that $\mathcal{V} = \mathbf{SP}(\mathbf{A})$. If a variety $\mathcal{V}$, like this, has a parallelogram term, then Theorem 6.16, combined with the main result of [8], proves that each quasivariety generator for $\mathcal{V}$ is dualizable.

Acknowledgment. The research reported in this paper was carried out while visiting La Trobe University as a guest of Brian Davey. We gratefully acknowledge the support and encouragement provided by our host. We also thank Ross Willard for reading a preliminary draft of the paper and recommending improvements.

REFERENCES

- [1] Berman, Joel; Idziak, Paweł; Marković, Petar; McKenzie, Ralph; Valeriote, Matthew; Willard, Ross *Varieties with few subalgebras of powers*. Trans. Amer. Math. Soc. **362** (2010), no. 3, 1445–1473.
- [2] Clark, David M.; Davey, Brian A. *Natural dualities for the working algebraist*. Cambridge Studies in Advanced Mathematics, **57**. Cambridge University Press, Cambridge, 1998.
- [3] Clark, David M.; Idziak, Paweł M.; Sabourin, Lousindi R.; Szabó, Csaba; Willard, Ross *Natural dualities for quasivarieties generated by a finite commutative ring*. Algebra Universalis **46** (2001), no. 1–2, 285–320.
- [4] Clark, David M.; Krauss, Peter H.; *Varieties generated by para primal algebras*. Algebra Universalis **7** (1977), no. 1, 93–114.
- [5] Davey, B. A.; Heindorf, L.; McKenzie, R. *Near unanimity: an obstacle to general duality theory*. Algebra Universalis **33** (1995), no. 3, 428–439.
- [6] Davey, B. A.; Quackenbush, R. W. *Bookkeeping duality for paraprimal algebras*. Contributions to general algebra, **9** (Linz, 1994), 19–26, Hölder-Pichler-Tempsky, Vienna, 1995.
- [7] Davey, B. A.; Werner, H. *Dualities and equivalences for varieties of algebras*. Contributions to lattice theory (Szeged, 1980), 101–275, Colloq. Math. Soc. János Bolyai, **33**, North-Holland, Amsterdam, 1983.
- [8] Davey, Brian A.; Willard, Ross *The dualisability of a quasi-variety is independent of the generating algebra*. Algebra Universalis **45** (2001), no. 1, 103–106.
- [9] Dent, Topaz; Kearnes, Keith A.; Szendrei, Ágnes *An easy test for congruence modularity*. (English summary) Algebra Universalis **67** (2012), no. 4, 375–392.
- [10] Freese, Ralph; McKenzie, Ralph *Commutator theory for congruence modular varieties*. London Mathematical Society Lecture Note Series, **125**. Cambridge University Press, Cambridge, 1987.

- [11] Idziak, Paweł *Congruence labelling for dualizable algebras*. unpublished notes, June 14, 1994.
- [12] Kearnes, Keith A *Residual bounds for varieties of modules*. Algebra Universalis **28** (1991), 448–452.
- [13] Kearnes, Keith A.; Szendrei, Ágnes *Clones of algebras with parallelogram terms*. Internat. J. Algebra Comput. **22** (2012), no. 1, 1250005, 30 pp.
- [14] McKenzie, Ralph *Residually small varieties of K -algebras*. Algebra Universalis **14** (1982), no. 2, 181–196.
- [15] McKenzie, Ralph *Narrowness implies uniformity*. Algebra Universalis **15** (1982), no. 1, 67–85.
- [16] Moore, Matthew *Naturally dualisable algebras omitting types 1 and 5 have a cube term*. <http://arxiv.org/abs/1403.7677>
- [17] Nickodemus, M. H. *Natural dualities for finite groups with abelian Sylow subgroups*. Thesis (Ph.D.)—University of Colorado at Boulder. 2007. 51 pp.
- [18] Pszczoła, Krzysztof J. *Duality for affine spaces over finite fields*. Contributions to general algebra, **13** (Velké Karlovice, 1999/Dresden, 2000), 285–293, Heyn, Klagenfurt, 2001.
- [19] Willard, Ross *Unpublished*.
- [20] Zádori, László *Natural duality via a finite set of relations*. Bull. Austral. Math. Soc. **51** (1995), no. 3, 469–478.

(Keith Kearnes) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COLORADO, BOULDER, CO 80309-0395, USA

E-mail address: Keith.Kearnes@Colorado.EDU

(Ágnes Szendrei) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COLORADO, BOULDER, CO 80309-0395, USA

E-mail address: Agnes.Szendrei@Colorado.EDU